

TCHP

Breast · NCCP 00722 · version 2

Your medicines

Your treatment schedule

A picture of side effects

Numbers connect each symptom box to the relevant anatomical area. Tell your team about new or worsening symptoms.

Trastuzumab / pertuzumab · HER2 treatment

1 Heart

Trastuzumab / pertuzumab

Heart function is monitored. Report new breathlessness, swelling or chest symptoms.

Docetaxel / carboplatin · chemotherapy

2 Bowel

Pertuzumab-containing treatment

Persistent or severe diarrhoea needs early advice.

3 Mouth / taste

Chemotherapy

Report mouth sores, difficulty drinking or significant taste changes.

4 Blood / infection

Docetaxel / carboplatin

Fever or sudden illness needs immediate oncology contact.

5 Nerves

Docetaxel

Report tingling or numbness, especially if persistent.

6 Skin / nails

Docetaxel-containing treatment

Report rash, nail changes or painful skin.

7 Infusion symptoms


Any infused medicine

Tell the nurse immediately about wheeze, flushing, rash, swelling or sudden symptoms during treatment.

8 Swelling

Treatment-related symptoms

New leg swelling or breathlessness needs assessment; symptoms may have several causes.



Selected effects only: the same symptom can have more than one cause, and not every possible side effect is shown.

TCHP

Breast · NCCP 00722 · version 2

TREATMENT JOURNEY

Before cycle 1 Baseline blood tests and cardiac assessment are reviewed.	Cycle day 1 Docetaxel + carboplatin + trastuzumab + pertuzumab are given.	Days 2-21 Recovery, supportive medicines and symptom monitoring.	Every 21 days The cycle repeats if review and blood tests allow.	After chemotherapy HER2-directed treatment usually continues according to the post-operative/adjuvant treatment plan.
--	---	--	--	---

IMPORTANT MEDICINE-SPECIFIC EFFECTS

Docetaxel - Severe neutropenia/infection can occur. - Fluid retention, nerve symptoms, nail/skin changes and infusion reactions can occur.	Carboplatin - Low blood counts, especially neutrophils and platelets, can occur. - Allergic infusion reactions can occur, particularly after repeated exposure. - Nausea and vomiting may occur despite preventative medicines.
Trastuzumab + pertuzumab - Heart function can fall and is monitored during HER2-directed treatment. - Infusion/administration reactions can occur. - Pertuzumab commonly causes diarrhoea; rare serious lung reactions can occur.	

OTHER / SHARED TREATMENT EFFECTS

The anatomy page highlights important symptom patterns for this exact regimen, but it cannot show every possible effect. Fatigue, nausea, appetite change, mouth/bowel symptoms or blood-count changes may occur depending on the medicines used. Report new or worsening symptoms even if they are not listed.

Monitoring / medicines - Follow the prescribed steroid premedication around docetaxel. - Growth-factor support may be part of the plan; follow the exact product and timing given by your oncology team. - Cardiac function is monitored because trastuzumab/pertuzumab can affect the heart.	Bring / track - Your current medicines, including non-prescription products. - A note of new symptoms, when they started and what they stopped you doing. - Any temperature, blood-pressure or bowel-count readings your team asked you to record. - Questions about the purpose of treatment, scan/review timing or changes to the plan.
--	--

WHEN TO CALL - DO NOT WAIT FOR THE NEXT APPOINTMENT

Call oncology urgently for fever, severe diarrhoea, new breathlessness, swelling, severe rash, uncontrolled vomiting, significant bleeding or a rapidly worsening symptom. Severe breathing difficulty or chest pain: call 112/999. If your team has given you different local thresholds or emergency instructions, follow those instructions.

Consent discussion support

Use the approved consent process to document treatment aim, expected benefit, material risks, alternatives, questions and the patient's decision.

Scan to open the regimen-specific patient portal

<https://sactcheck.com/patient/00722/>

Official NCCP 00722

SACTCheck v0.81.1 · patient information prototype for clinical review. Follow your oncology team and local emergency instructions.

