SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Gemcitabine alone: nasopharyngeal cancer

Head and neck: nasopharyngeal cancer · NCCP 00514 · version 3 · 21-day cycle

Patient name	DOB	Hospital no.	Date

Treatment context
Gemcitabine is used to control the cancer and relieve symptoms. Your team discusses expected benefit, alternatives and when to review whether treatment is helping.

Schedule
Gemcitabine 1,250 mg/m² on Days 1 and 8 every 21 days, for up to 6 cycles or until progression or unacceptable side effects.

1. TREATMENT DISCUSSION

- ☐ Purpose, expected benefit and alternatives explained
- ☐ Indication, medicines, sequence and duration confirmed
- ☐ Blood tests and any individually indicated heart checks discussed
- ☐ Support medicines and phase changes explained
- ☐ Fertility, contraception and breastfeeding discussed where relevant
- ☐ Common, serious and lasting drug-specific risks discussed below
- ☐ Urgent symptoms and local contact details confirmed

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Gemcitabine

ANTIMETABOLITE CHEMOTHERAPY

- ☐ Low blood counts, infection and bleeding
- ☐ Flu-like symptoms, fatigue, rash and swelling
- ☐ Uncommon serious lung or liver injury
- ☐ Rare haemolytic uraemic syndrome and kidney injury
- ☐ Rare severe skin reactions
- ☐ Effects on fertility; pregnancy and breastfeeding need specialist advice

Discuss treatment aim and alternatives, expected benefit and uncertainty, all listed drug risks, infection and emergency advice, blood and organ checks, support medicines, fertility and contraception, and the option of declining treatment. The linked NCCP source flags this nasopharyngeal gemcitabine use as outside the licensed indication; discuss and document consent under hospital policy.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00514 v3.

SACTCheck v0.84.49
Developed by Dr Paul O'Brien