National Cancer Control Programme (NCCP) 00514 · clinician draft

Gemcitabine alone: nasopharyngeal cancer

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Chemotherapy visit

Active medicines: Gemcitabine

Emetogenic risk: low

Medicine	Dose / route	When
Ondansetron	8 mg PO OD	Day 1 Clinician/pharmacy draft; reconcile to the individual prescription.
Metoclopramide	10 mg PO TDS	PRN Clinician/pharmacy draft; reconcile to the individual prescription.

Additional phase-specific support

- Gemcitabine is given into a vein over 30 minutes on Days 1 and 8. Checks, preparation and observation add to the time in the unit.
- Full blood count before each treatment; kidney and liver blood tests before each cycle.
- The team reviews symptoms, side effects and response. Treatment may finish earlier if it is no longer helping or causes unacceptable problems.
- The local CUH source is retained as a reference list, not as the national antiemetic standard. Nystatin/chlorhexidine are displayed as local mouth-care adjuncts, not universal antiemetic or mucositis prophylaxis. Verify current local oral-care policy.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00514 v3 · pTreatment / supportive care · NCCP regimen and NCIS antiemetic guidance

Sources checked 2026-11-11 · SACTCheck v0.84.49 · Regimen-level drafts; prescriber and oncology-pharmacy verification required