Gemcitabine 1250 and cisplatin: nasopharyngeal cancer

National Cancer Control Programme (NCCP) 00517 · Source reconciled draft; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Before your first visit

Full blood count, kidney and liver tests; review medicines, allergies, weight, oral intake and symptoms. Review nerve symptoms and hearing; audiology when indicated. Cisplatin eligibility includes adequate kidney function.

STEP 2

Time in the unit

Gemcitabine runs over 30 minutes, followed on Day 1 by cisplatin over 2 hours. Cisplatin requires prescribed fluids before and after treatment. Day 8 is gemcitabine alone.

STEP 3

Keep one calendar

Confirm the full treatment sequence and any separate radiotherapy plan. The linked NCCP document gives 4–6 cycles; this is not the three-cycle 00903 induction schedule.

STEP 4

Leaving the unit

Confirm the next tests, medicines, treatment dates and verified daytime and out-of-hours contact numbers.

STEP 5

Practical help

Ask about transport, work, costs, communication needs and whether a support person can attend.

Sources: NCCP 00517 v2: retrieved official regimen. Source reconciled draft; independent clinical, pharmacy and patient/PPI review pending.