Four-day carboplatin and fluorouracil with radiotherapy

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Days 1–4: four-day chemotherapy

Active medicines: Carboplatin + Fluorouracil

Emetogenic risk: moderate

Four consecutive carboplatin days: clinician/pharmacy must prescribe the complete multi-day antiemetic schedule. No unsupported fixed schedule is generated.

Additional phase-specific support

- Carboplatin runs over 30 minutes on each of Days 1–4. Fluorouracil runs continuously over 96 hours; the source also permits a separate 22-hour infusion on each of Days 1–4. Confirm the actual pump or bag-change and disconnection times.
- Before each cycle: full blood count, kidney and liver tests. Review weight, eating, drinking, swallowing, mouth symptoms, hands and feet. Kidney measurement and hearing tests follow the source and individual plan.
- Chemotherapy is given only during radiotherapy. If radiation is interrupted, contact the team before the next chemotherapy dose. Radiation oncology sets the radiation visits and follow-up.
- Preserve the CUH high/moderate and docetaxel proforma adjuncts where applicable: omeprazole, local nystatin/ chlorhexidine mouth care, breakthrough antiemetics and loperamide. These are historical local reference items requiring current-policy and patient-specific review, not universal prophylaxis or an additional duplicate prescription.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____