Carboplatin AUC 4 and fluorouracil with radiotherapy

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Chemotherapy visit

Active medicines: Carboplatin + Fluorouracil

Emetogenic risk: high

Medicine	Dose / route	When
Aprepitant	125 mg PO OD	Day 1 Clinician/pharmacy draft; reconcile to the individual prescription.
Ondansetron	16 mg PO OD	Day 1 Clinician/pharmacy draft; reconcile to the individual prescription.
Dexamethasone	12 mg PO OD	Day 1 Clinician/pharmacy draft; reconcile to the individual prescription.
Olanzapine	5 mg PO OD	Day 1 Clinician/pharmacy draft; reconcile to the individual prescription.
Aprepitant	80 mg PO OD	Days 2-3 Clinician/pharmacy draft; reconcile to the individual prescription.
Dexamethasone	8 mg PO OD	Days 2-4 Clinician/pharmacy draft; reconcile to the individual prescription.
Metoclopramide	10 mg PO TDS	Subsequent days / as directed Clinician/pharmacy draft; reconcile to the individual prescription.
Olanzapine	5 mg PO OD	Days 2-4 Clinician/pharmacy draft; reconcile to the individual prescription.

Additional phase-specific support

- Carboplatin is given over 30 minutes on Day 1; its dose is calculated from kidney function. Fluorouracil runs continuously over 96 hours; the source also permits separate 22-hour infusions on Days 1–4. Confirm the actual pump or bag-change and disconnection times.
- Before each cycle: full blood count, kidney and liver tests. Review weight, intake, swallowing, mouth symptoms, hands and feet. Measured kidney function is preferred when feasible; audiology is arranged when indicated.
- Chemotherapy is given only during radiotherapy. If radiation is interrupted, contact the team before the next chemotherapy dose. Radiation oncology sets the radiation visits and follow-up.
- The local CUH source is retained as a reference list, not as the national antiemetic standard. Nystatin/chlorhexidine are displayed as local mouth-care adjuncts, not universal antiemetic or mucositis prophylaxis. Verify current local oral-care policy.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____