Gemcitabine 1250 and cisplatin: nasopharyngeal cancer

The medicines act together to shrink or control nasopharyngeal cancer. Your team records how this course fits any planned radiation treatment and your overall treatment aim.


Head and neck: nasopharyngeal cancer · NCCP 00517 v2 · Document revised 11 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Confirm the full treatment sequence and any separate radiotherapy plan. The linked NCCP document gives 4–6 cycles; this is not the three-cycle 00903 induction schedule. Source check: the current HSE link supplies v2 while the NCIS list names v3. The treating team must reconcile this before clinical use.

My overall treatment

Day 1: both drugs; Day 8: gemcitabine. Every 21 days, 4–6 cycles.



Treatment
Gemcitabine 1,250 mg/m² on Days 1 and 8, with cisplatin 80 mg/m² on Day 1, every 21 days for 4–6 cycles or until progression or unacceptable side effects.

Review / next phase
Confirm the full treatment sequence and any separate radiotherapy plan. The linked NCCP document gives 4–6 cycles; this is not the three-cycle 00903 induction schedule.

A typical 21-day cycle

Day 1	Day 2–7	Day 8	Day 9–21
treatment	Between the first two visits	gemcitabine	Recovery and next review

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Baseline preparation** NCCP / individual plan	Full blood count, kidney and liver tests; review medicines, allergies, weight, oral intake and symptoms. Review nerve symptoms and hearing; audiology when indicated. Cisplatin eligibility includes adequate kidney function.
During	**Required tests** NCCP	Day 1: full blood count, kidney and liver tests. Day 8: full blood count and creatinine. Hearing assessment and measured kidney clearance are arranged when indicated.
After / response	**Recovery and next step** Individual plan	The team follows recovery, response and persistent side effects. Scans and tumour tests follow the individual plan; there is no universal scan or tumour-marker schedule in this regimen.

Between treatments

- Side effects can occur on any day. Keep prescribed support medicines and contact details to hand.
- Call promptly for fever or sudden illness, new breathing symptoms, persistent vomiting or inability to drink.

Bring your medicines list, appointment calendar and verified urgent contacts.

When to get help
Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.
Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321
EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.


1 Fatigue and concentration
Gemcitabine + Cisplatin
Tiredness can affect activity and concentration. Balance activity and rest; report sudden confusion.

3 Breathing
Gemcitabine + Cisplatin
New or worsening breathlessness, cough or chest symptoms need urgent advice; lung inflammation can be serious.

5 Sickness and bowels
Gemcitabine + Cisplatin
Take prescribed anti-sickness medicines. Report persistent vomiting, diarrhoea, dehydration or inability to drink.

7 Skin and infusion reactions
Gemcitabine + Cisplatin
Report severe rash, blistering, facial swelling or breathing symptoms promptly. Tell staff about pain at the infusion site.

9 Hearing and nerves
Cisplatin
Report ringing, hearing loss, numbness or tingling before the next dose; some effects can persist.



2 Mouth and swallowing
Gemcitabine + Cisplatin
Report soreness, ulcers or difficulty swallowing. Ask early about mouth care, pain relief and nutrition.

4 Blood and infection
Gemcitabine + Cisplatin
Fever, shivering, sudden illness, unexplained bruising or bleeding needs urgent oncology advice.

6 Kidneys and fluid balance
Gemcitabine + Cisplatin
Report reduced urine, swelling or poor fluid intake. Blood tests help detect kidney injury.

8 Fertility and pregnancy
Gemcitabine + Cisplatin
Discuss fertility preservation, contraception, pregnancy and breastfeeding before treatment.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Fever, shivering, sudden illness, unusual bleeding, new breathlessness, persistent vomiting or inability to drink needs urgent oncology advice. Severe chest pain, severe breathing difficulty, collapse or a seizure is an emergency: call 112/999.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Keep your treatment details to hand

Head and neck: nasopharyngeal cancer · NCCP 00517 v2 · Document revised 11 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Gemcitabine 1250 + cisplatin

Day 1: both drugs; Day 8: gemcitabine. Every 21 days, 4–6 cycles.

Phase: _____ Planned cycles: 6

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Team stamps after each completed cycle.

C1	C2	C3
Date / initials	Date / initials	Date / initials
C4	**C5**	**C6**
Date / initials	Date / initials	Date / initials

My treatment information

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.