Vismodegib for advanced basal cell skin cancer

NCCP 00236 · Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Taking or receiving treatment

Swallow the prescribed capsule whole with about 200 mL of water, with or without food, at the same time each day. Do not open, crush or chew it. Keep it in its labelled pack, away from children, and return unused capsules to the pharmacy.

STEP 2

Missed, vomited or held treatment

If a dose is missed, skip it and take the next dose at the usual time. Do not double up. If you vomit after taking it, ask the oncology team or pharmacist before taking more; do not assume that the dose needs replacing.

STEP 3

Appointments and supply

Arrange dates, medicine supply, transport and any needed interpreter or carer support. Ask whether bloods are taken in the oncology unit or elsewhere and how the result reaches the prescriber.

STEP 4

Paper copies and contact information

Keep this complete handbook, the three-page summary and the wallet card together. The QR code is optional. Ask the team to add your actual contacts and print a revised plan when treatment changes.

Sources: NCCP 00236 v3: current regimen source. Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending.