SACTCHECK · NCCP 00455

# My complete treatment handbook

## Pembrolizumab 200 mg · every 3 weeks · lung cancer

All information can be read on paper. Keep this with your treatment record.

> Use the urgent instructions and contact numbers in the following companion pages. Example telephone numbers must be replaced by your team before patient use. Your individual treatment plan takes precedence.

## Contents


1. Your three-page treatment companion — page 2
2. Full treatment reference: medicines, schedule, tests, symptoms and follow-up — page 4
3. Detailed treatment and monitoring plan — page 19
4. Supportive medicines and care — page 22
5. Visits, access and practical arrangements — page 24
6. Your treatment passport and symptom record — page 25
7. Wallet card and phone instructions — page 30


SACTCheck v0.84.40 · Clinical, pharmacy and patient review pending. Formal consent is supplied separately.

# Pembrolizumab 200 mg · every 3 weeks · lung cancer

Pembrolizumab 200 mg is given as a drip over about 30 minutes on Day 1 every 3 weeks (21 days). The hospital visit is longer because of checks and preparation. After complete surgery and platinum chemotherapy, the adjuvant course lasts up to 12 months. For metastatic cancer, treatment continues while it remains helpful and tolerable; the team may consider stopping after 24 months of sustained response without progression, or continuing when appropriate.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending. Selected effects; not a complete medicine leaflet.

## A picture of side effects

Numbered symptom boxes connect to the anatomical illustration. The medicine responsible may be uncertain; report the symptom.

### 1 Hormones / nervous system

**Pembrolizumab**

Severe headache, confusion, weakness, dizziness or marked thirst need prompt advice.

### 3 Heart

**Pembrolizumab**

Chest pain, racing heartbeat or fainting needs urgent assessment.

### 5 Kidneys


**Pembrolizumab**

Report much less urine, new swelling or a change in kidney tests.

### 7 Skin / muscles

**Pembrolizumab**

Spreading rash, blisters, severe weakness or muscle pain needs prompt review.




### 2 Lungs

**Pembrolizumab**

New cough or breathlessness can signal inflammation or another urgent cause.

### 4 Liver

**Pembrolizumab**

Yellow eyes/skin or dark urine may signal inflammation; tests also check the liver.

### 6 Bowel

**Pembrolizumab**

New or persistent diarrhoea, blood in stools or severe tummy pain needs urgent advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Contact the oncology team immediately for new or worsening breathlessness, chest pain, persistent diarrhoea, severe tummy pain, yellow skin/eyes, much less urine, severe headache, marked weakness, confusion, unusual thirst or passing much more urine, or a spreading/blistering rash. For severe breathing difficulty, collapse, seizure or other immediately dangerous symptoms, seek emergency care. Do not wait for the next infusion.

# Your treatment, at home.

## Important medicine-specific effects

### Pembrolizumab

- Immune inflammation of lungs, bowel, liver, kidneys, hormone glands, skin, heart and nervous system
- Infusion reactions
- Some effects need steroids, hospital care or long-term hormone replacement
- Effects may start after the final dose

**Surgery, radiation and monitoring:** open the whole-treatment map or scan the QR.

## Make the plan practical

### Sickness medicines

Pembrolizumab alone has a minimal risk of causing sickness. Routine anti-sickness medicines or infusion premedicines are not usually needed; your team can prescribe symptom relief if required.

### Immune alert card

Carry your alert card during treatment and afterwards. Show it when seeking urgent care; symptoms can be mistaken for an infection or an ordinary stomach upset.

### Hormones and steroids

Some immune effects need steroids or hormone replacement. Follow the prescribed plan and do not stop steroids suddenly. Ask for help if symptoms worsen or you cannot take essential medicines.

### Pregnancy and breastfeeding

Discuss pregnancy, fertility and breastfeeding before treatment. Use effective contraception during treatment and for at least four months after the last dose. If pregnancy is possible or occurs, contact your oncology team promptly.

**Emergency: 112 / 999.** Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

**Contact oncology immediately** for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

**Insert numbers here**

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

**Insert numbers here**

Ask for medical oncology

EVENINGS / OVERNIGHT

**Your out-of-hours pathway / nearest ED**

Other hospitals: use the numbers your team provides.

**Complete reference, visual timeline and passport**

https://sactcheck.com/patient/00455/

Sources and review status are available online. Follow your own prescription and hospital instructions.

NCCP 00455 · VERSION 16

# Your treatment, explained

Pembrolizumab 200 mg · every 3 weeks · lung cancer

21-day cycle | Source reconciled; independent clinical, pharmacy and patient/PPI review pending

ACT ON SYMPTOMS, NOT THE CALENDAR

## When to get help

**Emergency: call 112 / 999**

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

**Contact your oncology service immediately**

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Contact the oncology team immediately for new or worsening breathlessness, chest pain, persistent diarrhoea, severe tummy pain, yellow skin/eyes, much less urine, severe headache, marked weakness, confusion, unusual thirst or passing much more urine, or a spreading/blistering rash. For severe breathing difficulty, collapse, seizure or other immediately dangerous symptoms, seek emergency care. Do not wait for the next infusion.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.

Your treatment and urgent-contact card is supplied separately and in the wallet-card section of the complete handbook.

Sources: hse · nccp · smpc

YOUR COMPLETE REFERENCE

## Read what you need, when you need it.

Start with the overview, explore the cycle, or go straight to a question. Urgent advice is always available.

How each drug works
Your treatment and reviews
Your 21-day timeline
Baseline and cycle-specific checks
Whole treatment, scans and checks
Tablets, visits and practical planning
Your lung cancer and treatment aim
How pembrolizumab works
Your treatment course
Tests and reviews
Practical support and everyday life
When to get urgent help
Supportive medicines
Understanding the evidence
Questions to ask
Sources

UNDERSTAND YOUR TREATMENT


# How pembrolizumab works

For metastatic lung cancer, the aim is to control cancer and help people live longer. After surgery and platinum chemotherapy, the adjuvant aim is to reduce the chance of cancer returning. Ask which aim applies to you.

IMMUNOTHERAPY

## Pembrolizumab

Keytruda



### Blocks an immune brake

Pembrolizumab is an antibody that blocks PD-1, a control switch on immune T cells. Cancer cells can use this switch to reduce immune attack. Blocking it can help the immune system recognise and attack cancer. The same increased immune activity can inflame healthy organs, which is why new symptoms need attention even after treatment ends.

Immune cell → PD-1 blocked → Cancer attacked

**What to watch for**

Report new breathing, bowel, skin, hormone, heart or nerve symptoms promptly; these may be immune inflammation.

Sources: smpc

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

The anatomical side-effect map follows.

THE BIGGER TREATMENT PLAN

# Your treatment and reviews

ONE IMMUNOTHERAPY MEDICINE

## Pembrolizumab every 3 weeks

C1 C2 C3 …

After complete surgery and platinum chemotherapy, the adjuvant course lasts up to 12 months. For metastatic cancer, treatment continues while it remains helpful and tolerable; the team may consider stopping after 24 months of sustained response without progression, or continuing when appropriate.

Your team confirms your treatment aim, duration and review plan. Treatment may change after discussion.

Sources: nccp

BASELINE → TREATMENT → FOLLOW-UP

# Checks across your treatment

This is a relative treatment map, not a booking calendar. Ask who books each test and reviews the result. Extra tests may be needed for symptoms.

| Check | Before starting | First infusion | Between infusions | Course review | After treatment |
|---|---|---|---|---|---|
| **Blood count, kidney, liver and glucose** | Baseline blood tests | Results reviewed before dosing | Before every 3-week cycle | Confirm cause of abnormal tests; clinician decides hold, restart or stop | Follow unresolved abnormalities |
| **Thyroid and other hormones** | Baseline thyroid function; other hormones if needed | Review symptoms and hormone medicines | Thyroid every 3–6 weeks; cortisol if clinically indicated | Confirm replacement and steroid-taper plan | Continue hormone follow-up if needed |
| **Symptoms and immune effects** | Discuss existing illnesses and warning symptoms | Tell the nurse immediately about a reaction | Report new symptoms on any day | Hold/restart decisions need clinical review | Late immune effects remain possible |
| **Cancer response and course length** | Confirm metastatic or adjuvant plan | Agree how benefit will be assessed | Scans and review on your team's schedule | Adjuvant: up to 12 months; metastatic: individual 24-month review | Agree surveillance or next treatment |

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00455 v16 · official regimen

THE WHOLE TREATMENT JOURNEY

# Pembrolizumab 200 mg · every 3 weeks · lung cancer: treatment and checks

### Lung cancer only: first-line metastatic NSCLC or adjuvant NSCLC after surgery and platinum chemotherapy

Pembrolizumab 200 mg is given as a drip over about 30 minutes on Day 1 every 3 weeks (21 days). The hospital visit is longer because of checks and preparation. After complete surgery and platinum chemotherapy, the adjuvant course lasts up to 12 months. For metastatic cancer, treatment continues while it remains helpful and tolerable; the team may consider stopping after 24 months of sustained response without progression, or continuing when appropriate.

**Read from left to right.** Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

**Treatment, tests and reviews across the course**

| Care / check | Before starting | First infusion | Between infusions | Course review | After treatment |
|---|---|---|---|---|---|
| **Treatment** NCCP 00455 | Confirm metastatic or adjuvant indication | 200 mg IV over about 30 minutes | Day1 every 3 weeks while prescribed | Adjuvant maximum 12 months; metastatic individual review | Follow-up/surveillance or next treatment |
| **Blood count, kidney, liver and glucose** NCCP 00455 | Baseline blood tests | Results reviewed before dosing | Before every 3-week cycle | Confirm cause of abnormal tests; clinician decides hold, restart or stop | Follow unresolved abnormalities |
| **Thyroid and other hormones** NCCP 00455 · SMPC | Baseline thyroid function; other hormones if needed | Review symptoms and hormone medicines | Thyroid every 3–6 weeks; cortisol if clinically indicated | Confirm replacement and steroid-taper plan | Continue hormone follow-up if needed |
| **Symptoms and immune effects** NCCP 00455 · SMPC | Discuss existing illnesses and warning symptoms | Tell the nurse immediately about a reaction | Report new symptoms on any day | Hold/restart decisions need clinical review | Late immune effects remain possible |
| **Cancer response and course length** NCCP 00455 | Confirm metastatic or adjuvant plan | Agree how benefit will be assessed | Scans and review on your team's schedule | Adjuvant: up to 12 months; metastatic: individual 24-month review | Agree surveillance or next treatment |

## What each check is for

1

### Before first dose

Before starting

Confirm the indication and baseline safety

**Ask / act:** Confirm results, access, consent discussion, alert card and responsible teams.

Sources: NCCP 00455

2

### Course-length decision

At planned course review

Metastatic and adjuvant durations differ

**Ask / act:** Document whether treatment ends, pauses or continues, and how follow-up is arranged.

Sources: NCCP 00455

3

**After the last dose**

After treatment

Immune effects and hormone needs can continue

**Ask / act:** Provide the monitoring and urgent-contact plan.

Sources: NCCP 00455 · SMPC

4

**Abnormal liver or kidney results**

Whenever a result changes

A blood result alone does not establish immune inflammation

**Ask / act:** Contact the treating team and confirm the investigation and treatment plan.

Sources: NCCP 00455 · SMPC

## At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-10-29 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.26

### Day 1 → Day 21: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

Pembrolizumab 200 mg is given as a drip over about 30 minutes on Day 1 every 3 weeks (21 days). The hospital visit is longer because of checks and preparation. After complete surgery and platinum chemotherapy, the adjuvant course lasts up to 12 months. For metastatic cancer, treatment continues while it remains helpful and tolerable; the team may consider stopping after 24 months of sustained response without progression, or continuing when appropriate.

**Treatment timing: displayed days 1–21**

| Treatment or check | Days in this window | What happens |
|---|---|---|
| **Infusion** | Day 1 | Day 1 · hospital infusion |
| **Immune effects** | Days 1–21 | Any day · also after treatment ends |
| **Energy, skin and bowels** | Days 1–21 | Timing varies · report changes promptly |

**Symptoms can occur at any time:** Treatment days and symptom timing are different: there is no reliably safe week for immune effects.

**Confirm your treatment aim and duration:** After complete surgery and platinum chemotherapy, the adjuvant course lasts up to 12 months. For metastatic cancer, treatment continues while it remains helpful and tolerable; the team may consider stopping after 24 months of sustained response without progression, or continuing when appropriate.

**Why these bands stay broad:** The next Day 1 is normally 21 days after this cycle's Day 1, if your team agrees it is safe to proceed. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment.

**There are no guaranteed "safe days".** Immune effects can happen on any day and after treatment ends. Carry your immunotherapy alert card and tell any healthcare team that you receive pembrolizumab.

### Complete timing and symptom notes

#### Day 1: Your infusion visit

The team reviews your blood results, symptoms and medicines. Tell the nurse straight away about itching, flushing, a tight chest, dizziness or feeling suddenly unwell during the drip. Before leaving, confirm the next tests, treatment visit and urgent contact number.

#### Day 2–21: Between visits

You may feel well between infusions, but immune inflammation can begin at any time. Keep a short symptom record and take medicines exactly as prescribed. Contact your team about new symptoms instead of waiting for your next appointment. Confirm thyroid tests are booked every 3–6 weeks, even if infusions are six weeks apart.

PRACTICAL TREATMENT GUIDE

## Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

### Appointments

Pembrolizumab 200 mg is given as a drip over about 30 minutes on Day 1 every 3 weeks (21 days). The hospital visit is longer because of checks and preparation.

STEP 2

### Course length

After complete surgery and platinum chemotherapy, the adjuvant course lasts up to 12 months. For metastatic cancer, treatment continues while it remains helpful and tolerable; the team may consider stopping after 24 months of sustained response without progression, or continuing when appropriate.

STEP 3

### Checks between visits

Blood tests are due before each infusion. Thyroid tests stay every 3–6 weeks regardless of dose schedule. Ask about travel, work, transport and whom to contact if a test or visit is delayed.

STEP 4

### Home medicines and records

Keep the alert card, symptom record, current medicine list and urgent contact details accessible. Your team may prescribe steroids or hormone replacement only if needed.

STEP 5

### After the final dose

Keep reporting new symptoms. Some hormone medicines and monitoring continue after the cancer treatment finishes.

Sources: NCCP 00455 v16 · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.

MAKE THE PLAN PRACTICAL

## Support around the medicines

### 1 Sickness medicines

Pembrolizumab alone has a minimal risk of causing sickness. Routine anti-sickness medicines or infusion premedicines are not usually needed; your team can prescribe symptom relief if required.

### 2 Immune alert card

Carry your alert card during treatment and afterwards. Show it when seeking urgent care; symptoms can be mistaken for an infection or an ordinary stomach upset.

### 3 Hormones and steroids

Some immune effects need steroids or hormone replacement. Follow the prescribed plan and do not stop steroids suddenly. Ask for help if symptoms worsen or you cannot take essential medicines. If treatment is paused because of an immune side effect, your team checks recovery and any steroid or hormone plan before deciding whether it can restart. Do not change steroids or restart treatment yourself.

### 4 Pregnancy and breastfeeding

Discuss pregnancy, fertility and breastfeeding before treatment. Use effective contraception during treatment and for at least four months after the last dose. If pregnancy is possible or occurs, contact your oncology team promptly.

Sources: nccp · smpc

TREATMENT REFERENCE · 01

## Your lung cancer and treatment aim

First confirm which lung setting applies to you.

### Metastatic treatment

Metastatic means the cancer has spread. First-line means the first medicine treatment for this stage. Here, pembrolizumab alone is offered for suitable cancers with a high PD-L1 result and without EGFR or ALK alterations. The aim is usually control, symptom relief and more time.

Sources: nccp · smpc

### After surgery

Adjuvant means treatment after surgery to reduce the chance of cancer returning. This pathway follows complete resection and platinum chemotherapy for cancer at high risk of recurrence. It does not use the same high PD-L1 requirement as the metastatic pathway.

Sources: nccp · smpc

### Your alternatives

Ask which other treatments, clinical trials, surveillance or symptom-focused care fit your exact cancer and health. Supportive care can help during either pathway.

Sources: nccp · smpc

TREATMENT REFERENCE · 02

## How pembrolizumab works

This is an immunotherapy medicine.

### Helping immune cells recognise cancer

Pembrolizumab is an antibody that blocks PD-1, a control switch on immune T cells. Cancer cells can use this switch to reduce immune attack. Blocking it can help the immune system recognise and attack cancer. The same increased immune activity can inflame healthy organs, which is why new symptoms need attention even after treatment ends.

Sources: nccp · smpc

### Why healthy organs can be affected

The immune system can also attack healthy tissues. Lung, bowel, liver, kidney, skin, gland, heart and nerve inflammation can range from mild to serious. Some hormone changes need long-term replacement.

Sources: nccp · smpc

### A pause is a safety decision

Your team may delay a dose, treat side effects or stop pembrolizumab. The usual response to immune toxicity is to hold or stop treatment, not to give a smaller dose.

Sources: nccp · smpc

TREATMENT REFERENCE · 03

## Your treatment course

The appointment pattern and total duration are different decisions.

### Infusion schedule

Pembrolizumab 200 mg is given as a drip over about 30 minutes on Day 1 every 3 weeks (21 days). The hospital visit is longer because of checks and preparation.

Sources: nccp · smpc

### Course length

After complete surgery and platinum chemotherapy, the adjuvant course lasts up to 12 months. For metastatic cancer, treatment continues while it remains helpful and tolerable; the team may consider stopping after 24 months of sustained response without progression, or continuing when appropriate.

Sources: nccp · smpc

### Changing schedules

These lung trials used pembrolizumab 200 mg every 3 weeks. Trial treatment limits and research scan schedules are not automatic NCCP prescribing rules.

Sources: nccp · smpc

TREATMENT REFERENCE · 04

## Tests and reviews

Tell your team about symptoms even when recent blood tests were normal.

### Before starting

Your team checks the cancer diagnosis, treatment aim and relevant tumour tests. Blood count, kidney/liver, glucose and thyroid tests provide a baseline. Discuss existing illnesses, medicines, pregnancy and breastfeeding.

Sources: nccp · smpc

### During treatment

Blood count, kidney/liver function and glucose before every 3-week cycle; thyroid every 3–6 weeks, cortisol if indicated, and symptom review throughout and after treatment.

Sources: nccp · smpc

### During a pause or after stopping

Extra tests may be needed for symptoms. Follow the agreed steroid or hormone plan. Keep your alert card and continue reporting symptoms after the last dose.

Sources: nccp · smpc

TREATMENT REFERENCE · 05

## Practical support and everyday life

Keep your team informed about what matters to you.

### Energy and activity

Fatigue can affect daily activities. Balance activity with rest and ask for help if tiredness is new, marked or accompanied by dizziness, headache, thirst or weakness; it may need medical assessment.

Sources: nccp · smpc

### Eating, drinking and bowels

Drink as advised and report ongoing vomiting or diarrhoea. Do not assume diarrhoea is a minor stomach upset; your team needs to check for immune inflammation and other causes.

Sources: nccp · smpc

### Medicines and pregnancy

Bring a full medicine list, including supplements and steroids. Ask before starting new medicines. Use the contraception plan and discuss fertility or breastfeeding with your team.

Sources: nccp · smpc

TREATMENT REFERENCE · 06

## When to get urgent help

Use the contact plan given by your oncology service.

### Report warning symptoms

Contact the oncology team immediately for new or worsening breathlessness, chest pain, persistent diarrhoea, severe tummy pain, yellow skin/eyes, much less urine, severe headache, marked weakness, confusion, unusual thirst or passing much more urine, or a spreading/blistering rash. For severe breathing difficulty, collapse, seizure or other immediately dangerous symptoms, seek emergency care. Do not wait for the next infusion.

Sources: nccp · smpc

### During the drip

Tell the nurse immediately if you become itchy, flushed, breathless, dizzy, tight-chested or suddenly unwell.

Sources: nccp · smpc

### Care after treatment ends

Immune inflammation can start late. Tell any healthcare team about pembrolizumab even after the course has finished.

Sources: nccp · smpc

## Why this treatment may be offered

In a trial of suitable people with metastatic lung cancer and a high PD-L1 result, pembrolizumab helped people live longer than chemotherapy. After surgery and chemotherapy, a separate trial found a lower risk of recurrence or death. Its final long-term analysis did not demonstrate a statistically significant improvement in overall survival. The aims and evidence differ between these settings.

These results describe groups of people, not what will happen to one person. The metastatic and after-surgery studies answer different questions.

Sources: keynote024 · keynote091 · smpc · keynote091-final

PREPARE FOR THE CONVERSATION

## Questions worth asking

### Is my treatment for metastatic cancer or to reduce recurrence after surgery?

Ask your team to write down the treatment aim. For metastatic disease, the aim is usually cancer control, symptom relief and more time. After complete surgery and platinum chemotherapy, adjuvant treatment aims to reduce the chance of recurrence.

### Which tumour-test results support this plan?

For the first-line metastatic monotherapy pathway, the source requires high PD-L1 expression and no EGFR mutation or ALK translocation. The adjuvant pathway has different eligibility. Ask which results apply to your own cancer.

### How long is my course, and what could change it?

The adjuvant course lasts up to 12 months. Metastatic treatment continues while helpful and tolerable; a 24-month review may lead to stopping or continuing. Symptoms, response and your preferences can change the plan.

**Who arranges blood and thyroid tests between visits?**

The oncology team arranges and reviews tests. Blood count, kidney/liver function and glucose are checked before each cycle; thyroid tests are usually every 3–6 weeks. Ask whether an extra test is needed between six-weekly infusions.

**Which symptoms should I report straight away, and whom do I contact out of hours?**

Report new or worsening breathing problems, persistent diarrhoea, severe abdominal pain, yellow eyes, reduced urine, severe headache, weakness, thirst or a spreading rash promptly. Keep the local oncology and out-of-hours numbers available; severe breathing difficulty, collapse, seizure or severe chest pain needs 112 / 999.

**What are my alternatives if I choose not to have this treatment?**

Ask about alternatives for your exact stage and tumour results, including other anticancer treatments, trials, surveillance where appropriate and symptom-focused care. Supportive care remains available whether or not you proceed.

**Does an abnormal blood test mean pembrolizumab must be stopped permanently?**

Not by itself. Your team checks whether the change is caused by immune inflammation, another medicine, cancer or another illness. They decide whether to pause treatment, investigate further or stop it. Do not change the treatment plan yourself.

> **Try explaining the plan back:** Which medicines are you having now? What changes after induction? What would make you call today? Ask your team to fill any gaps.

SOURCE TRANSPARENCY

## Where this information comes from

Regimen schedule: NCCP 00455 v16. Medicine explanations and safety information: sources below. The timeline is an educational illustration, not a measured risk curve or an individual prediction.


- NCCP 00455 v16 · official regimen
- EMA Keytruda IV product information and patient leaflet
- KEYNOTE-024 five-year randomized follow-up · Reck et al. 2021
- PEARLS/KEYNOTE-091 randomized adjuvant study · O’Brien et al. 2022
- Lala et al. 2020 · 400 mg six-week exposure-bridging study (with corrigendum)
- HSE · chemotherapy safety and when to contact your team
- EORTC primary trial-sponsor report · PEARLS/KEYNOTE-091 final analysis, ESMO 2026

## Written source addresses

https://healthservice.hse.ie/documents/6903/Pembrolizumab_200mg_Monotherapy_455.pdf

https://www.ema.europa.eu/en/documents/product-information/keytruda-epar-product-information_en.pdf

https://pubmed.ncbi.nlm.nih.gov/33872070/

https://pubmed.ncbi.nlm.nih.gov/36108662/

https://pubmed.ncbi.nlm.nih.gov/32305010/

https://www2.hse.ie/conditions/chemotherapy/side-effects/

https://www.eortc.org/blog/2026/10/27/eortc-at-esmo-2026/

BASELINE → TREATMENT → FOLLOW-UP

# Checks across your treatment

This is a relative treatment map, not a booking calendar. Ask who books each test and reviews the result. Extra tests may be needed for symptoms.

| Check | Before starting | First infusion | Between infusions | Course review | After treatment |
|---|---|---|---|---|---|
| **Blood count, kidney, liver and glucose** | Baseline blood tests | Results reviewed before dosing | Before every 3-week cycle | Confirm cause of abnormal tests; clinician decides hold, restart or stop | Follow unresolved abnormalities |
| **Thyroid and other hormones** | Baseline thyroid function; other hormones if needed | Review symptoms and hormone medicines | Thyroid every 3–6 weeks; cortisol if clinically indicated | Confirm replacement and steroid-taper plan | Continue hormone follow-up if needed |
| **Symptoms and immune effects** | Discuss existing illnesses and warning symptoms | Tell the nurse immediately about a reaction | Report new symptoms on any day | Hold/restart decisions need clinical review | Late immune effects remain possible |
| **Cancer response and course length** | Confirm metastatic or adjuvant plan | Agree how benefit will be assessed | Scans and review on your team’s schedule | Adjuvant: up to 12 months; metastatic: individual 24-month review | Agree surveillance or next treatment |

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00455 v16 · official regimen

THE WHOLE TREATMENT JOURNEY

# Pembrolizumab 200 mg · every 3 weeks · lung cancer: treatment and checks

**Lung cancer only: first-line metastatic NSCLC or adjuvant NSCLC after surgery and platinum chemotherapy**

Pembrolizumab 200 mg is given as a drip over about 30 minutes on Day 1 every 3 weeks (21 days). The hospital visit is longer because of checks and preparation. After complete surgery and platinum chemotherapy, the adjuvant course lasts up to 12 months. For metastatic cancer, treatment continues while it remains helpful and tolerable; the team may consider stopping after 24 months of sustained response without progression, or continuing when appropriate.

**Read from left to right.** Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

**Treatment, tests and reviews across the course**

| Care / check | Before starting | First infusion | Between infusions | Course review | After treatment |
|---|---|---|---|---|---|
| **Treatment** NCCP 00455 | Confirm metastatic or adjuvant indication | 200 mg IV over about 30 minutes | Day1 every 3 weeks while prescribed | Adjuvant maximum 12 months; metastatic individual review | Follow-up/surveillance or next treatment |
| **Blood count, kidney, liver and glucose** NCCP 00455 | Baseline blood tests | Results reviewed before dosing | Before every 3-week cycle | Confirm cause of abnormal tests; clinician decides hold, restart or stop | Follow unresolved abnormalities |
| **Thyroid and other hormones** NCCP 00455 · SMPC | Baseline thyroid function; other hormones if needed | Review symptoms and hormone medicines | Thyroid every 3–6 weeks; cortisol if clinically indicated | Confirm replacement and steroid-taper plan | Continue hormone follow-up if needed |
| **Symptoms and immune effects** NCCP 00455 · SMPC | Discuss existing illnesses and warning symptoms | Tell the nurse immediately about a reaction | Report new symptoms on any day | Hold/restart decisions need clinical review | Late immune effects remain possible |
| **Cancer response and course length** NCCP 00455 | Confirm metastatic or adjuvant plan | Agree how benefit will be assessed | Scans and review on your team's schedule | Adjuvant: up to 12 months; metastatic: individual 24-month review | Agree surveillance or next treatment |

# What each check is for

### 1 Before first dose

Before starting

Confirm the indication and baseline safety

**Ask / act:** Confirm results, access, consent discussion, alert card and responsible teams.

Sources: NCCP 00455

### 2 Course-length decision

At planned course review

Metastatic and adjuvant durations differ

**Ask / act:** Document whether treatment ends, pauses or continues, and how follow-up is arranged.

Sources: NCCP 00455

### 3 After the last dose

After treatment

Immune effects and hormone needs can continue

**Ask / act:** Provide the monitoring and urgent-contact plan.

Sources: NCCP 00455 · SMPC

### 4 Abnormal liver or kidney results

Whenever a result changes

A blood result alone does not establish immune inflammation

**Ask / act:** Contact the treating team and confirm the investigation and treatment plan.

Sources: NCCP 00455 · SMPC

## At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-10-29 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.26

# Pembrolizumab 200 mg · every 3 weeks · lung cancer

**Lung cancer information** This patient suite covers the two lung indications described here. The shared clinical protocol also lists other cancers, which need their own patient information.

**Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

## Separate scripts for each phase

Choose the active phase in the clinician workflow. Each script has its own medicines, timing and review checks.

- **Pembrolizumab 200 mg every 3 weeks** — Minimal pembrolizumab

https://sactcheck.com/patient/00455/supportive-care.html

# Pembrolizumab 200 mg · every 3 weeks · lung cancer

**Lung cancer information** This patient suite covers the two lung indications described here. The shared clinical protocol also lists other cancers, which need their own patient information.

**Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

## Pembrolizumab 200 mg every 3 weeks

**Active medicines:** pembrolizumab

**Emetogenic risk:** Minimal

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

### Additional phase-specific support

- No routine antiemetic prophylaxis or routine infusion premedication is usually required. Use an individual/local symptom-relief prescription if needed.
- Carry the immunotherapy alert card and report possible immune symptoms promptly during and after treatment.
- Blood count, renal/liver profile and glucose before each cycle; thyroid every 3–6 weeks, cortisol if clinically indicated.
- Effective contraception during treatment and at least 4 months after the last dose.
- If treatment is paused because of an immune side effect, your team checks recovery and any steroid or hormone plan before deciding whether it can restart. Do not change steroids or restart treatment yourself.

**Before signing:** verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00455 v16 · pNCCP00455 pp5–7 · NCCP00455 v16 supportive care

Sources checked 2026-10-29 · SACTCheck v0.84.26 · Regimen-level drafts; prescriber and oncology-pharmacy verification required

# Pembrolizumab 200 mg · every 3 weeks · lung cancer

NCCP 00455 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

## Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

### Appointments

Pembrolizumab 200 mg is given as a drip over about 30 minutes on Day 1 every 3 weeks (21 days). The hospital visit is longer because of checks and preparation.

STEP 2

### Course length

After complete surgery and platinum chemotherapy, the adjuvant course lasts up to 12 months. For metastatic cancer, treatment continues while it remains helpful and tolerable; the team may consider stopping after 24 months of sustained response without progression, or continuing when appropriate.

STEP 3

### Checks between visits

Blood tests are due before each infusion. Thyroid tests stay every 3–6 weeks on either dose schedule. Ask about travel, work, transport and whom to contact if a test or visit is delayed.

STEP 4

### Home medicines and records

Keep the alert card, symptom record, current medicine list and urgent contact details accessible. Your team may prescribe steroids or hormone replacement only if needed.

STEP 5

### After the final dose

Keep reporting new symptoms. Some hormone medicines and monitoring continue after the cancer treatment finishes.

Sources: NCCP 00455 v16 · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.

# Pembrolizumab 200 mg · every 3 weeks · lung cancer

## A record of what you experience

One cycle per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal CTCAE grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Pembrolizumab 200 mg is given as a drip over about 30 minutes on Day 1 every 3 weeks (21 days). The hospital visit is longer because of checks and preparation. After complete surgery and platinum chemotherapy, the adjuvant course lasts up to 12 months. For metastatic cancer, treatment continues while it remains helpful and tolerable; the team may consider stopping after 24 months of sustained response without progression, or continuing when appropriate.

The four pages are reusable recording sheets, not a prescribed four-cycle course. Ask for additional sheets as needed.

**Emergency: 112 / 999.** Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

**Contact oncology immediately** for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

**Insert numbers here**

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

**Insert numbers here**

Ask for medical oncology

EVENINGS / OVERNIGHT

**Your out-of-hours pathway / nearest ED**

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

If treatment is paused because of an immune side effect, your team checks recovery and any steroid or hormone plan before deciding whether it can restart. Do not change steroids or restart treatment yourself.

https://sactcheck.com/patient/00455/

Full patient information, sources and visual timeline. No identifiers are encoded.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

# Pembrolizumab 200 mg · every 3 weeks · lung cancer

Cycle number: _____ Start date: _____ Next review: _____

**3-week monotherapy cycle 1 · confirm tests and symptoms** · copy this sheet for another cycle of the same phase.

### Pembrolizumab · Immunotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Breathing / cough** | ☐ Usual breathing | ☐ New cough or breathlessness: call promptly | ☐ Symptoms with usual activity: urgent review | ☐ Breathless at rest: urgent assessment | ☐ Severe breathing difficulty / collapse: 112/999 |
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | ☐ None | ☐ Mild change: report | ☐ Pain or daily tasks affected: call | ☐ Spreading infection or blistering: urgent review | ☐ Severe illness / collapse: emergency help |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Pembrolizumab 200 mg · every 3 weeks · lung cancer

Cycle number: _____ Start date: _____ Next review: _____

**3-week monotherapy cycle 2 · confirm tests and symptoms** · copy this sheet for another cycle of the same phase.

### Pembrolizumab · Immunotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Breathing / cough** | ☐ Usual breathing | ☐ New cough or breathlessness: call promptly | ☐ Symptoms with usual activity: urgent review | ☐ Breathless at rest: urgent assessment | ☐ Severe breathing difficulty / collapse: 112/999 |
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | ☐ None | ☐ Mild change: report | ☐ Pain or daily tasks affected: call | ☐ Spreading infection or blistering: urgent review | ☐ Severe illness / collapse: emergency help |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Pembrolizumab 200 mg · every 3 weeks · lung cancer

Cycle number: _____ Start date: _____ Next review: _____

**3-week monotherapy cycle 3 · confirm tests and symptoms** · copy this sheet for another cycle of the same phase.

### Pembrolizumab · Immunotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Breathing / cough** | ☐ Usual breathing | ☐ New cough or breathlessness: call promptly | ☐ Symptoms with usual activity: urgent review | ☐ Breathless at rest: urgent assessment | ☐ Severe breathing difficulty / collapse: 112/999 |
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | ☐ None | ☐ Mild change: report | ☐ Pain or daily tasks affected: call | ☐ Spreading infection or blistering: urgent review | ☐ Severe illness / collapse: emergency help |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Pembrolizumab 200 mg · every 3 weeks · lung cancer

Cycle number: _____ Start date: _____ Next review: _____

**3-week monotherapy cycle 4 · confirm tests and symptoms** · copy this sheet for another cycle of the same phase.

### Pembrolizumab · Immunotherapy

| Symptom | None | Mild | Moderate / call | Severe / urgent | Emergency signs |
|---|---|---|---|---|---|
| **Breathing / cough** | ☐ Usual breathing | ☐ New cough or breathlessness: call promptly | ☐ Symptoms with usual activity: urgent review | ☐ Breathless at rest: urgent assessment | ☐ Severe breathing difficulty / collapse: 112/999 |
| **Diarrhoea** | ☐ Usual pattern | ☐ Up to 3 extra stools above normal | ☐ 4–6 extra stools/day or at night: call | ☐ 7+ extra stools/day or dehydration: urgent review | ☐ Collapse / severe dehydration: 112/999 |
| **Skin / nail changes** | ☐ None | ☐ Mild change: report | ☐ Pain or daily tasks affected: call | ☐ Spreading infection or blistering: urgent review | ☐ Severe illness / collapse: emergency help |
| **Fatigue** | ☐ Usual energy | ☐ Tiredness improves with rest | ☐ Rest does not help; usual tasks affected: call | ☐ Self-care difficult: urgent review | ☐ Collapse: 112/999 |

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

# Carry your treatment details

Print, cut out and fold. Follow the contact plan from your own hospital.

SACTCHECK · SACT TREATMENT

## Pembrolizumab 200 mg · every 3 weeks · lung cancer

NCCP 00455 v16 · 21-day cycle

Pembrolizumab 200 mg is given as a drip over about 30 minutes on Day 1 every 3 weeks (21 days). The hospital visit is longer because of checks and preparation. After complete surgery and platinum chemotherapy, the adjuvant course lasts up to 12 months. For metastatic cancer, treatment continues while it remains helpful and tolerable; the team may consider stopping after 24 months of sustained response without progression, or continuing when appropriate.

| 1 | 2 | 3 | 4 | 5 | 6 |
|---|---|---|---|---|---|

**Your contact plan**

Oncology team: Insert numbers here
Out of hours: Insert numbers here.

Evenings / overnight: instructed out-of-hours pathway or nearest ED.
**Emergency: 112 / 999.**

Other hospital / own contact: _____

QR: https://sactcheck.com/patient/00455/ · no patient information.

Draft for review · SACTCheck v0.84.26