SACTCHECK · NCCP 00812

My complete treatment handbook

Cemiplimab for advanced cutaneous squamous cell cancer

All information can be read on paper. Keep this with your treatment record.

> Use the urgent instructions and contact numbers in the following companion pages. Example telephone numbers must be replaced by your team before patient use. Your individual treatment plan takes precedence.

Contents

1. Your three-page treatment companion — page 2
2. Full treatment reference: medicines, schedule, tests, symptoms and follow-up — page 5
3. Detailed treatment and monitoring plan — page 24
4. Supportive medicines and care — page 27
5. Visits, access and practical arrangements — page 29
6. Your treatment passport and symptom record — page 30
7. Wallet card and phone instructions — page 34

SACTCheck v0.84.40 · Clinical, pharmacy and patient review pending. Formal consent is supplied separately.

Cemiplimab for advanced cutaneous squamous cell cancer

Immunotherapy for advanced skin SCC when curative surgery or radiotherapy is not appropriate.

NCCP 00812b: metastatic or locally advanced cutaneous squamous cell carcinoma when curative surgery or curative radiotherapy is not appropriate. This guide covers skin SCC, not the separate cervical-cancer indication. · NCCP 00812 v3a · Document revised 6 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

NCCP 00812b: metastatic or locally advanced cutaneous squamous cell carcinoma when curative surgery or curative radiotherapy is not appropriate. This guide covers skin SCC, not the separate cervical-cancer indication. This information supports individual consent; it does not issue or change a prescription.

My overall treatment

Cemiplimab infusion every 3 weeks; no fixed cycle total in this regimen.

Cemiplimab treatment
Continue while beneficial and tolerable, with team review; no fixed cycle total.

A typical 21-day cycle

Day 1	Day 2–21
Treatment visit	Between visits

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Baseline safety** NCCP / medicine-specific precautions	Blood count, kidneys/liver, glucose, thyroid and hepatitis B/C tests.
During	**Bloods and organ checks** NCCP; local interval clarification where stated	Blood count, kidneys/liver and glucose each cycle; thyroid TSH every 3–6 weeks.
After / response	**Follow-up** Regimen / individual disease plan	Continue reporting immune symptoms after stopping; scans, skin and lesion reviews as planned.

Between treatments

- Immune symptoms can occur between visits or after the last dose.
- Tell every clinician about immunotherapy and carry the alert card.

Bring the written medicine plan, test dates and actual oncology/out-of-hours contacts.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321
EXAMPLE ONLY — team must replace both numbers before patient use.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Hormones / head / eyes

Cemiplimab

New severe headache, visual change, unusual tiredness, dizziness, thirst or frequent urination may signal immune effects and needs prompt advice.

2 Lungs / breathing

Cemiplimab

New cough or breathlessness may reflect lung inflammation or another serious cause. Call promptly; severe breathing difficulty is an emergency.

3 Heart / muscles


Cemiplimab

Chest pain, palpitations, marked muscle weakness or swallowing difficulty can be serious immune effects. Seek urgent assessment.

4 Liver

Cemiplimab

Inflammation may first show in blood tests. Report yellow eyes or skin, dark urine, unusual itching or upper-abdominal pain.



5 Kidneys / urine

Cemiplimab

Kidney inflammation may be detected on blood tests. Report reduced urine, swelling, blood in urine or a marked change from normal.

6 Bowels / abdomen

Cemiplimab

Report increased stool frequency, diarrhoea, blood or mucus, or abdominal pain early. Severe pain, black stools or dehydration needs urgent help.

7 Skin / mouth

Cemiplimab

Report rash or itching early. Blistering, peeling, mouth or genital sores, or rash with fever needs urgent assessment.

8 Nerves / strength

Cemiplimab

New tingling, weakness, unsteadiness, facial change or trouble swallowing needs prompt review. Sudden weakness or speech difficulty is an emergency.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Immunotherapy can cause inflammation in healthy organs during treatment or after it stops. Contact the oncology team promptly for new diarrhoea, blood in stools, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell any clinician that you are receiving or have received immunotherapy.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Keep your treatment details to hand

NCCP 00812b: metastatic or locally advanced cutaneous squamous cell carcinoma when curative surgery or curative radiotherapy is not appropriate. This guide covers skin SCC, not the separate cervical-cancer indication. · NCCP 00812 v3a · Document revised 6 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Cemiplimab

Cemiplimab infusion every 3 weeks; no fixed cycle total in this regimen.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Use an ongoing dated visit record; there is no preset completion stamp total.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

NCCP 00812 · VERSION 3A

Your treatment, explained

NCCP 00812b: metastatic or locally advanced cutaneous squamous cell carcinoma when curative surgery or curative radiotherapy is not appropriate. This guide covers skin SCC, not the separate cervical-cancer indication. · NCCP 00812 v3a · Document revised 6 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Cemiplimab for advanced cutaneous squamous cell cancer

Reference treatment cycle

Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

ACT ON SYMPTOMS, NOT THE CALENDAR

When to get help

Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Immunotherapy can cause inflammation in healthy organs during treatment or after it stops. Contact the oncology team promptly for new diarrhoea, blood in stools, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell any clinician that you are receiving or have received immunotherapy.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.

Your treatment and urgent-contact card is supplied separately and in the wallet-card section of the complete handbook.

Sources: nccp

YOUR COMPLETE REFERENCE

Read what you need, when you need it.

Start with the overview, explore the cycle, or go straight to a question. Urgent advice is always available.

How each drug works
The overall treatment plan
Your 21-day timeline
Baseline and cycle-specific checks
Whole treatment, scans and checks
Tablets, visits and practical planning
Understanding your diagnosis and treatment goal
Your medicines and the overall plan
Taking treatment and preparing for visits
Tests: what is required and what each result means
What to expect between visits
Skin checks, sun protection and optional photographs
Pregnancy, fertility and protecting others
Questions, decisions and practical support
Follow-up and your continuing record
Supportive medicines
Understanding the evidence
Questions to ask
Sources

UNDERSTAND YOUR TREATMENT


Medicines in this regimen

Cemiplimab helps immune cells act against advanced cutaneous squamous cell cancer when curative surgery or radiotherapy is not suitable. The team reviews cancer control, symptoms, side effects and your priorities to decide whether to continue.

PD-1 CHECKPOINT IMMUNOTHERAPY

Cemiplimab

Libtayo



Releases a PD-1 immune brake

Cemiplimab blocks PD-1 signalling that can limit T-cell activity. This may help the immune system attack cutaneous squamous cell cancer. Increased immune activity can also inflame healthy organs, so new symptoms need attention even if the visible skin tumour appears to improve.

PD-1 blocked → Immune response activated → Organ inflammation possible

What to watch for

Carry an immunotherapy alert card and report new diarrhoea, cough, breathlessness, rash, weakness or endocrine symptoms promptly.

Sources: nccp · nccn-patient

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

The anatomical side-effect map follows.

REGIMEN-SPECIFIC SIDE EFFECTS

Your anatomical side-effect map

A picture of side effects

Match each symptom box to its numbered body location. Tell your team about new symptoms.

1 Hormones / head / eyes

Cemiplimab

New severe headache, visual change, unusual tiredness, dizziness, thirst or frequent urination may signal immune effects and needs prompt advice.

3 Heart / muscles

Cemiplimab

Chest pain, palpitations, marked muscle weakness or swallowing difficulty can be serious immune effects. Seek urgent assessment.

5 Kidneys / urine


Cemiplimab

Kidney inflammation may be detected on blood tests. Report reduced urine, swelling, blood in urine or a marked change from normal.

7 Skin / mouth

Cemiplimab

Report rash or itching early. Blistering, peeling, mouth or genital sores, or rash with fever needs urgent assessment.



2 Lungs / breathing

Cemiplimab

New cough or breathlessness may reflect lung inflammation or another serious cause. Call promptly; severe breathing difficulty is an emergency.

4 Liver

Cemiplimab

Inflammation may first show in blood tests. Report yellow eyes or skin, dark urine, unusual itching or upper-abdominal pain.

6 Bowels / abdomen

Cemiplimab

Report increased stool frequency, diarrhoea, blood or mucus, or abdominal pain early. Severe pain, black stools or dehydration needs urgent help.

8 Nerves / strength

Cemiplimab

New tingling, weakness, unsteadiness, facial change or trouble swallowing needs prompt review. Sudden weakness or speech difficulty is an emergency.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

THE BIGGER TREATMENT PLAN

The overall treatment plan

EVERY THREE WEEKS

Cemiplimab treatment

No preset cycle total **Cemiplimab infusion every 3 weeks; no fixed cycle total in this regimen.**

Cemiplimab is given by IV infusion on Day 1 every 21 days. The usual source dose is 350 mg over 30 minutes; the prescription and visit checks confirm your treatment. There is no fixed cycle total in this regimen: treatment continues while the team considers it beneficial and tolerable.

The treatment setting determines duration; a reference calendar is not a prescription or a reason to restart a held medicine.

Sources: nccp

BASELINE → TREATMENT → FOLLOW-UP

Required safety checks and your individual disease-assessment plan

The team supplies actual dates and resolves any source ambiguity. A reference period does not replace a prescription, a local monitoring plan or an individual treatment decision.

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00812 v3a: current regimen source · NCCN Guidelines for Patients: Squamous Cell Skin Cancer, 2023 - supplied education reference

THE WHOLE TREATMENT JOURNEY

Cemiplimab for advanced cutaneous squamous cell cancer: treatment and checks

NCCP 00812b: metastatic or locally advanced cutaneous squamous cell carcinoma when curative surgery or curative radiotherapy is not appropriate. This guide covers skin SCC, not the separate cervical-cancer indication.

Cemiplimab is given by IV infusion on Day 1 every 21 days. The usual source dose is 350 mg over 30 minutes; the prescription and visit checks confirm your treatment. There is no fixed cycle total in this regimen: treatment continues while the team considers it beneficial and tolerable.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	During treatment	Between visits	Response / follow-up	After a hold or stop
Bloods and organ safety NCCP 00812	Full blood count, kidney/liver tests, glucose and thyroid tests; hepatitis B surface antigen, hepatitis B core antibody and hepatitis C testing. Review autoimmune conditions, organ transplantation, previous immune reactions and medicines.	Full blood count, kidney/liver tests and glucose each cycle; thyroid-stimulating hormone (TSH) every three to six weeks. Review new symptoms at every visit and arrange additional tests when indicated. Response assessment and scans follow the individual plan.	Contact the team for new symptoms; extra tests may be needed between scheduled checks.	Safety bloods do not by themselves show whether cancer is responding.	Immune effects can begin or persist after treatment stops. Keep the agreed follow-up, steroid or hormone-replacement instructions and alert card, and tell clinicians about previous cemiplimab. The next scan, skin/lesion review and any further treatment are individually planned.
Cancer, skin and daily function NCCP 00812	Record diagnosis, stage, current disease sites and the baseline assessment.	Keep the source-specific skin and clinical reviews and the individually agreed response assessments.	Record new lesions, lumps, wound changes, pain and effects on usual activities.	Write the next examination or scan date and the responsible team; no universal scan or tumour-marker interval is assumed.	Follow-up includes examination of the treated area, the rest of the skin and relevant lymph nodes. Imaging and specialist visits depend on the extent of disease, risk features, symptoms and treatment response. Keep a clear wound and skin-surveillance plan alongside oncology visits.
Medicines and continuing precautions NCCP 00812	Review all medicines and the pregnancy/ fertility plan.	Review treatment taken or received, holds, adverse effects and supportive medicines.	Keep the actual urgent contacts and do not restart a held medicine yourself.	Bring the medicine list and treatment summary to other services.	Use effective contraception during cemiplimab treatment and for at least four months after the last dose. Discuss fertility before treatment, and follow the medicine-specific breastfeeding restrictions. Report a possible pregnancy promptly.

What each check is for

1

Cemiplimab treatment

Every three weeks

Cemiplimab helps immune cells act against advanced cutaneous squamous cell cancer when curative surgery or radiotherapy is not suitable. The team reviews cancer control, symptoms, side effects and your priorities to decide whether to continue.

Ask / act: Confirm the current treatment phase, prescription, tests and support medicines.

Sources: NCCP 00812

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-06 · Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.40

Day 1 → Day 21: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

Cemiplimab is given by IV infusion on Day 1 every 21 days. The usual source dose is 350 mg over 30 minutes; the prescription and visit checks confirm your treatment. There is no fixed cycle total in this regimen: treatment continues while the team considers it beneficial and tolerable.

Treatment timing: displayed days 1–21

Treatment or check	Days in this window	What happens
Cemiplimab infusion	Day 1	Treatment after clinical review and the required tests.
Symptoms between visits	Days 1–21	Immune effects can begin at any time, including after treatment stops.
Next-cycle assessment	Day 21	The team confirms whether the next dose can proceed.

Timing note 1: Cemiplimab is given by IV infusion on Day 1 every 21 days. The usual source dose is 350 mg over 30 minutes; the prescription and visit checks confirm your treatment. There is no fixed cycle total in this regimen: treatment continues while the team considers it beneficial and tolerable.

Timing note 2: If a visit is missed or treatment is held, contact the unit for a new appointment and treatment decision. Do not assume the next scheduled date means treatment has been cleared to restart.

Timing note 3: This patient suite covers cutaneous SCC only; it does not transfer cervical-cancer treatment context.

Timing note 4: The cited NCCP indication is advanced disease unsuitable for curative surgery/radiotherapy; no adjuvant CSCC claim is made.

Timing note 5: A visible skin response does not exclude toxicity or establish response elsewhere.

Why these bands stay broad: This calendar shows the pattern. Your team supplies actual dates, prescribed doses and any hold or restart instructions. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00812 v3a: current regimen source ↗️

There are no guaranteed "safe days". The prescribed dose, treatment decision and local urgent-contact plan take priority. Call 112 or 999 for severe breathing difficulty, severe chest pain, collapse, a new seizure, sudden weakness or speech difficulty, or uncontrolled bleeding. Follow your oncology unit's written urgent-contact plan for other new or worsening symptoms; do not wait for the next appointment.

Complete timing and symptom notes

Day 1: Treatment visit

Cemiplimab is infused into a vein over 30 minutes in the source schedule. Allow additional time for assessment, preparation and observation. Tell staff immediately about itching, flushing, chills, pain, dizziness, breathlessness or swelling during the infusion. Report symptoms and previous reactions before treatment.

Day 2–21: Between visits

Immunotherapy can cause inflammation in healthy organs during treatment or after it stops. Contact the oncology team promptly for new diarrhoea, blood in stools, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell any clinician that you are receiving or have received immunotherapy. Do not wait for the next infusion to report a possible immune effect.

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Taking or receiving treatment

Cemiplimab is infused into a vein over 30 minutes in the source schedule. Allow additional time for assessment, preparation and observation. Tell staff immediately about itching, flushing, chills, pain, dizziness, breathlessness or swelling during the infusion.

STEP 2

Missed, vomited or held treatment

If a visit is missed or treatment is held, contact the unit for a new appointment and treatment decision. Do not assume the next scheduled date means treatment has been cleared to restart.

STEP 3

Appointments and supply

Arrange dates, medicine supply, transport and any needed interpreter or carer support. Ask whether bloods are taken in the oncology unit or elsewhere and how the result reaches the prescriber.

STEP 4

Paper copies and contact information

Keep this complete handbook, the three-page summary and the wallet card together. The QR code is optional. Ask the team to add your actual contacts and print a revised plan when treatment changes.

Sources: NCCP 00812 v3a: current regimen source. Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending.

MAKE THE PLAN PRACTICAL

Support around the medicines

1 An individual symptom plan

Ask which anti-sickness or other symptom medicines are prescribed, when to take them and when to call if they do not help. Do not start a borrowed medicine or alter treatment to manage a side effect yourself.

2 Your urgent contacts

Keep your actual oncology, out-of-hours and pharmacy contacts on the wallet card. Example numbers on a template must be replaced before you use it.

3 Pregnancy and fertility precautions

Use effective contraception during cemiplimab treatment and for at least four months after the last dose. Discuss fertility before treatment, and follow the medicine-specific breastfeeding restrictions. Report a possible pregnancy promptly.

4 Eating, activity and practical help

Report falling intake, weight loss, pain or fatigue early. The team can tailor nutrition, activity, wound, psychological and practical support to your needs.

5 Steroids and hormone replacement

Some immune effects need steroids or lasting hormone replacement. Follow the exact prescription and taper. Carry the relevant alert information, and do not stop or reduce these medicines yourself when anticancer treatment is held or completed.

Sources: nccp · nccn-patient

TREATMENT REFERENCE · 01

Understanding your diagnosis and treatment goal

Use these questions with your own pathology report and agreed treatment plan.

Confirm that this is skin SCC

This guide concerns cutaneous squamous cell carcinoma arising in the skin. Squamous cancers beginning inside the mouth, throat, lung or cervix have different treatment pathways. Record the original skin site, pathology date and whether the cancer has returned or spread.

Sources: nccn-patient

Why a cancer is considered high risk

The size, depth, site, growth pattern and involvement of nerves, lymph nodes or other structures can affect the plan. Prior treatment, recurrence and reduced immunity also matter. Ask the team to explain the risk features in your report in ordinary language; a risk category is not itself a treatment response.

Sources: nccn-patient

Why medicine is being offered

The cited NCCP cemiplimab setting is advanced skin SCC when curative surgery or radiotherapy is not appropriate. Record the multidisciplinary team's reason and the aim of treatment. This guide does not make a general promise that immunotherapy will avoid surgery or cure the cancer.

Sources: nccn-patient

Immunity and transplantation

Tell the oncology team about an organ transplant, immune-suppressing medicines, autoimmune illness and previous immune treatment. These can substantially change the balance of benefit and harm. Never stop transplant or steroid medicines yourself; the relevant specialists need a shared plan.

Sources: nccn-patient

Lesions, wounds and daily function

A skin tumour or treatment can affect pain, bleeding, wound care, movement, eating, appearance or sleep. Ask for a wound and symptom plan that names who to contact. Worsening bleeding, infection symptoms, new nerve pain or weakness should be assessed rather than waiting for a photograph comparison.

Sources: nccn-patient

TREATMENT REFERENCE · 02

Your medicines and the overall plan

Cemiplimab helps immune cells act against advanced cutaneous squamous cell cancer when curative surgery or radiotherapy is not suitable. The team reviews cancer control, symptoms, side effects and your priorities to decide whether to continue.

The schedule in this setting

Cemiplimab is given by IV infusion on Day 1 every 21 days. The usual source dose is 350 mg over 30 minutes; the prescription and visit checks confirm your treatment. There is no fixed cycle total in this regimen: treatment continues while the team considers it beneficial and tolerable.

Sources: nccp

How the treatment works

Cemiplimab blocks PD-1 signalling that can limit T-cell activity. This may help the immune system attack cutaneous squamous cell cancer. Increased immune activity can also inflame healthy organs, so new symptoms need attention even if the visible skin tumour appears to improve.

Sources: nccp

The role of earlier or later treatment

Record surgery, radiotherapy and other anticancer medicines separately, including dates and the specialist responsible. This schedule does not automatically include or authorise another treatment. Ask whether any overlap changes tests, symptoms to report or the timing of appointments.

Sources: nccn-patient

Pauses and stopping

A safety pause is an active treatment decision. Record which medicine is paused, the reason, the tests or recovery needed and who may authorise restarting. Do not restart from an old prescription or reset the history when a new phase begins.

Sources: nccp

TREATMENT REFERENCE · 03

Taking treatment and preparing for visits

Keep the written medicine plan beside this guide; it records your own dose and dates.

Taking or receiving the medicine

Cemiplimab is infused into a vein over 30 minutes in the source schedule. Allow additional time for assessment, preparation and observation. Tell staff immediately about itching, flushing, chills, pain, dizziness, breathlessness or swelling during the infusion.

Sources: nccp

Missed, vomited or held treatment

If a visit is missed or treatment is held, contact the unit for a new appointment and treatment decision. Do not assume the next scheduled date means treatment has been cleared to restart.

Sources: nccp

Other medicines and supplements

Show the pharmacist a current list of prescriptions, over-the-counter medicines, vitamins, herbal products and supplements. Include anticoagulants, heart medicines, steroids and recent antibiotics. Check before adding, stopping or changing anything; an apparently harmless product can alter treatment or confuse a symptom assessment.

Sources: nccp

Practical preparation

Arrange medicine supply, transport, interpreter or carer support before it is needed. Bring the alert card, symptom record and questions. Ask which appointments require someone to accompany you and how to obtain urgent advice outside normal hours.

Sources: nccn-patient

TREATMENT REFERENCE · 04

Tests: what is required and what each result means

Safety tests, tumour tests and response scans answer different questions.

Before treatment

Full blood count, kidney/liver tests, glucose and thyroid tests; hepatitis B surface antigen, hepatitis B core antibody and hepatitis C testing. Review autoimmune conditions, organ transplantation, previous immune reactions and medicines.

Sources: nccp

During treatment

Full blood count, kidney/liver tests and glucose each cycle; thyroid-stimulating hormone (TSH) every three to six weeks. Review new symptoms at every visit and arrange additional tests when indicated. Response assessment and scans follow the individual plan.

Sources: nccp

Scans and tumour markers

Ask the team to write the next assessment and who will arrange it. Bloods used to monitor kidneys, liver, hormones or muscle are not automatically measures of cancer response. There is no universal tumour-marker blood schedule for these skin-cancer regimens; a scan or extra blood test is requested for a specific clinical reason.

Sources: nccp

After a pause or finish

Immune effects can begin or persist after treatment stops. Keep the agreed follow-up, steroid or hormone-replacement instructions and alert card, and tell clinicians about previous cemiplimab. The next scan, skin/lesion review and any further treatment are individually planned.

Sources: nccp

Getting the result

For each test, record its purpose, due date, result and action. Agree how results reach you and who checks a result from another hospital. If an appointment or result has not arrived, contact the named service rather than assuming no news means no action is needed.

Sources: nccn-patient

TREATMENT REFERENCE · 05

What to expect between visits

You can feel well and still need the planned checks. New symptoms deserve attention even on a day without an appointment.

Your usual pattern

Record the start and change in symptoms, temperature if advised, bowel pattern, drinking, urine and the effect on normal tasks. A simple few-line record is enough. Use it to describe a change rather than trying to grade or diagnose a toxicity yourself.

Sources: nccp

When to call

Immunotherapy can cause inflammation in healthy organs during treatment or after it stops. Contact the oncology team promptly for new diarrhoea, blood in stools, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell any clinician that you are receiving or have received immunotherapy.

Sources: nccp

Eating, drinking and activity

Tell the team early about nausea, taste change, a sore mouth, diarrhoea or falling weight. Ask for an individual fluid and food plan if heart, kidney or other conditions limit intake. Gentle activity and planned rest can help maintain function; adjust to symptoms and ask for support when fatigue limits daily life.

Sources: nccn-patient

A held dose and a new phase

If a visit is missed or treatment is held, contact the unit for a new appointment and treatment decision. Do not assume the next scheduled date means treatment has been cleared to restart.

Sources: nccp

TREATMENT REFERENCE · 06

Skin checks, sun protection and optional photographs

This complements clinical assessment and does not replace it.

Protect your skin

Use shade, protective clothing, a hat and broad-spectrum sunscreen of at least SPF 30, and avoid tanning beds. Reapply sunscreen according to its instructions. Protection matters even when it is cloudy; choose a practical approach that lets you remain active safely.

Sources: nccn-patient

Know the pattern of your skin

Ask how to examine your skin, scars and relevant lymph-node areas and how often your own team recommends doing so. A mirror or a trusted helper can make hard-to-see areas easier to check. Report a new or changing lesion, a non-healing sore or a new lump rather than waiting for a routine visit.

Sources: nccn-patient

Photographs are optional

If useful, keep a dated photograph with consistent lighting and a size reference, following your team's advice. Photographs are optional and do not diagnose a cancer or replace examination. Keep them privately; this printable passport does not require you to upload images or other personal health information.

Sources: nccn-patient

Wounds, nails and sensitive skin

Use the dressings and skin-care plan recommended for your lesion and treatment. Ask before applying new creams or remedies to an open or treated area. Worsening pain, discharge, redness, bleeding or loss of function needs advice; routine skin-care instructions are not a substitute for assessment.

Sources: nccn-patient

TREATMENT REFERENCE · 07

Pregnancy, fertility and protecting others

Discuss these precautions before treatment even if pregnancy is not currently planned.

Medicine-specific precautions

Use effective contraception during cemiplimab treatment and for at least four months after the last dose. Discuss fertility before treatment, and follow the medicine-specific breastfeeding restrictions. Report a possible pregnancy promptly.

Sources: nccp

Planning ahead

Ask early about fertility preservation, sexual health and symptoms affecting intimacy. Tell the team about any chance of pregnancy, breastfeeding or a partner's pregnancy so that the correct medicine-specific advice can be given. Keep the precautions and their end dates in the written plan.

Sources: nccp

Safe handling and supply

Do not share anticancer medicines. Keep oral medicines in their original labelled packaging, away from children and anyone who could take them accidentally. Return unused medicines to the supplying pharmacy and follow its handling advice. A treatment hold does not remove pregnancy precautions.

Sources: nccp

TREATMENT REFERENCE · 08

Questions, decisions and practical support

Your priorities and the burden of treatment belong in the discussion.

Benefits and uncertainties

Ask what evidence applies to your diagnosis, stage, biomarker and treatment setting, what outcome the team hopes to improve, and what may happen without this treatment. A trial average cannot predict an individual result; a benefit measured in one setting should not be transferred to another.

Sources: nccn-patient

Alternatives and review points

Discuss other suitable treatments, clinical trials if relevant, local treatment options and symptom-focused care. Ask when the decision will be reviewed and what would prompt a change. Supportive and palliative care can help with symptoms and quality of life alongside anticancer treatment.

Sources: nccn-patient

Help with daily life

Tell the team about work, transport, caring responsibilities, cost concerns, body image, anxiety or sleep. Ask which nurse, social worker, dietitian, rehabilitation professional or support organisation can help. Use resources relevant to your country and service; a US guide may describe different access or funding.

Sources: nccn-patient

Shared understanding

Ask someone on the team to check that you can explain the medicine schedule, next tests and urgent-contact route in your own words. An interpreter, large-print copy or a trusted supporter can help; none requires you to use a smartphone.

Sources: nccn-patient

TREATMENT REFERENCE · 09

Follow-up and your continuing record

Keep this record when treatment is paused, changed or completed.

Your follow-up plan

Follow-up includes examination of the treated area, the rest of the skin and relevant lymph nodes. Imaging and specialist visits depend on the extent of disease, risk features, symptoms and treatment response. Keep a clear wound and skin-surveillance plan alongside oncology visits.

Sources: nccn-patient

Continuing medicine precautions

Immune effects can begin or persist after treatment stops. Keep the agreed follow-up, steroid or hormone-replacement instructions and alert card, and tell clinicians about previous cemiplimab. The next scan, skin/lesion review and any further treatment are individually planned.

Sources: nccp

Treatment summary for other clinicians

Record the medicines and last dates, important reactions, any long-term hormone replacement or steroid taper, and the responsible oncology contact. Show this information in urgent care and to other specialists. A different treatment phase does not erase earlier toxicity.

Sources: nccp

The complete paper version

This handbook can be used without a phone: keep the schedule, symptom advice, medicine list, tests and contacts together. The QR code is an optional route to the matching regimen information. Ask the team to print updates when your plan changes and keep the latest dated copy.

Sources: nccn-patient

What the evidence means for your treatment

The evidence discussion must match the diagnosis, biomarker and treatment setting shown in this guide.

Ask which outcomes matter in your setting and what benefit is realistic for you. No survival percentage from another regimen, dose combination or disease setting is assigned here.

Sources: nccp · nccn-patient

PREPARE FOR THE CONVERSATION

Questions worth asking

What is the aim of my treatment?

Cemiplimab helps immune cells act against advanced cutaneous squamous cell cancer when curative surgery or radiotherapy is not suitable. The team reviews cancer control, symptoms, side effects and your priorities to decide whether to continue.

Is there a fixed number of cycles?

Cemiplimab is given by IV infusion on Day 1 every 21 days. The usual source dose is 350 mg over 30 minutes; the prescription and visit checks confirm your treatment. There is no fixed cycle total in this regimen: treatment continues while the team considers it beneficial and tolerable.

What happens if treatment is held?

If a visit is missed or treatment is held, contact the unit for a new appointment and treatment decision. Do not assume the next scheduled date means treatment has been cleared to restart.

When are my next tests?

Full blood count, kidney/liver tests and glucose each cycle; thyroid-stimulating hormone (TSH) every three to six weeks. Review new symptoms at every visit and arrange additional tests when indicated. Response assessment and scans follow the individual plan. Your team adds the actual dates.

Can I use this without a smartphone?

Yes. The full handbook, three-page summary and wallet card are printable. The QR code is optional; keep actual contacts and written appointments on the paper copy.

> **Try explaining the plan back:** Can you explain your treatment setting, medicine schedule, next tests, the symptoms that need urgent advice and the number you would call?

SOURCE TRANSPARENCY

Where this information comes from

Regimen schedule: NCCP 00812 v3a. Medicine explanations and safety information: sources below. The timeline is an educational illustration, not a measured risk curve or an individual prediction.

- NCCP 00812 v3a: current regimen source
- NCCN Guidelines for Patients: Squamous Cell Skin Cancer, 2023 - supplied education reference

SACTCheck · Developed by Dr Paul O'Brien · Sources checked 2026-11-06.
This information supports your treatment discussion. Your oncology team confirms your personal treatment and urgent-care plan.

Written source addresses

https://healthservice.hse.ie/documents/6426/812_V3a_Cemiplimab.pdf

https://www.nccn.org/patients/guidelines/content/PDF/squamous_cell-patient.pdf

BASELINE → TREATMENT → FOLLOW-UP

Required safety checks and your individual disease-assessment plan

The team supplies actual dates and resolves any source ambiguity. A reference period does not replace a prescription, a local monitoring plan or an individual treatment decision.

Check	Before starting	During treatment	Between visits	Response / follow-up	After a hold or stop
Bloods and organ safety	Full blood count, kidney/liver tests, glucose and thyroid tests; hepatitis B surface antigen, hepatitis B core antibody and hepatitis C testing. Review autoimmune conditions, organ transplantation, previous immune reactions and medicines.	Full blood count, kidney/liver tests and glucose each cycle; thyroid-stimulating hormone (TSH) every three to six weeks. Review new symptoms at every visit and arrange additional tests when indicated. Response assessment and scans follow the individual plan.	Contact the team for new symptoms; extra tests may be needed between scheduled checks.	Safety bloods do not by themselves show whether cancer is responding.	Immune effects can begin or persist after treatment stops. Keep the agreed follow-up, steroid or hormone-replacement instructions and alert card, and tell clinicians about previous cemiplimab. The next scan, skin/lesion review and any further treatment are individually planned.
Cancer, skin and daily function	Record diagnosis, stage, current disease sites and the baseline assessment.	Keep the source-specific skin and clinical reviews and the individually agreed response assessments.	Record new lesions, lumps, wound changes, pain and effects on usual activities.	Write the next examination or scan date and the responsible team; no universal scan or tumour-marker interval is assumed.	Follow-up includes examination of the treated area, the rest of the skin and relevant lymph nodes. Imaging and specialist visits depend on the extent of disease, risk features, symptoms and treatment response. Keep a clear wound and skin-surveillance plan alongside oncology visits.
Medicines and continuing precautions	Review all medicines and the pregnancy/fertility plan.	Review treatment taken or received, holds, adverse effects and supportive medicines.	Keep the actual urgent contacts and do not restart a held medicine yourself.	Bring the medicine list and treatment summary to other services.	Use effective contraception during cemiplimab treatment and for at least four months after the last dose. Discuss fertility before treatment, and follow the medicine-specific breastfeeding restrictions. Report a possible pregnancy promptly.

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00812 v3a: current regimen source · NCCN Guidelines for Patients: Squamous Cell Skin Cancer, 2023 - supplied education reference

THE WHOLE TREATMENT JOURNEY

Cemiplimab for advanced cutaneous squamous cell cancer: treatment and checks

NCCP 00812b: metastatic or locally advanced cutaneous squamous cell carcinoma when curative surgery or curative radiotherapy is not appropriate. This guide covers skin SCC, not the separate cervical-cancer indication.

Cemiplimab is given by IV infusion on Day 1 every 21 days. The usual source dose is 350 mg over 30 minutes; the prescription and visit checks confirm your treatment. There is no fixed cycle total in this regimen: treatment continues while the team considers it beneficial and tolerable.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	During treatment	Between visits	Response / follow-up	After a hold or stop
Bloods and organ safety NCCP 00812	Full blood count, kidney/liver tests, glucose and thyroid tests; hepatitis B surface antigen, hepatitis B core antibody and hepatitis C testing. Review autoimmune conditions, organ transplantation, previous immune reactions and medicines.	Full blood count, kidney/liver tests and glucose each cycle; thyroid-stimulating hormone (TSH) every three to six weeks. Review new symptoms at every visit and arrange additional tests when indicated. Response assessment and scans follow the individual plan.	Contact the team for new symptoms; extra tests may be needed between scheduled checks.	Safety bloods do not by themselves show whether cancer is responding.	Immune effects can begin or persist after treatment stops. Keep the agreed follow-up, steroid or hormone-replacement instructions and alert card, and tell clinicians about previous cemiplimab. The next scan, skin/lesion review and any further treatment are individually planned.
Cancer, skin and daily function NCCP 00812	Record diagnosis, stage, current disease sites and the baseline assessment.	Keep the source-specific skin and clinical reviews and the individually agreed response assessments.	Record new lesions, lumps, wound changes, pain and effects on usual activities.	Write the next examination or scan date and the responsible team; no universal scan or tumour-marker interval is assumed.	Follow-up includes examination of the treated area, the rest of the skin and relevant lymph nodes. Imaging and specialist visits depend on the extent of disease, risk features, symptoms and treatment response. Keep a clear wound and skin-surveillance plan alongside oncology visits.
Medicines and continuing precautions NCCP 00812	Review all medicines and the pregnancy/fertility plan.	Review treatment taken or received, holds, adverse effects and supportive medicines.	Keep the actual urgent contacts and do not restart a held medicine yourself.	Bring the medicine list and treatment summary to other services.	Use effective contraception during cemiplimab treatment and for at least four months after the last dose. Discuss fertility before treatment, and follow the medicine-specific breastfeeding restrictions. Report a possible pregnancy promptly.

What each check is for

1

Cemiplimab treatment

Every three weeks

Cemiplimab helps immune cells act against advanced cutaneous squamous cell cancer when curative surgery or radiotherapy is not suitable. The team reviews cancer control, symptoms, side effects and your priorities to decide whether to continue.

Ask / act: Confirm the current treatment phase, prescription, tests and support medicines.

Sources: NCCP 00812

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-06 · Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.40

Cemiplimab for advanced cutaneous squamous cell cancer

Your diagnosis and treatment setting NCCP 00812b: metastatic or locally advanced cutaneous squamous cell carcinoma when curative surgery or curative radiotherapy is not appropriate. This guide covers skin SCC, not the separate cervical-cancer indication.

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Separate scripts for each phase

Choose the active phase in the clinician workflow. Each script has its own medicines, timing and review checks.

- **Cemiplimab treatment** — Minimal
  Cemiplimab

https://sactcheck.com/patient/00812/supportive-care.html

Cemiplimab for advanced cutaneous squamous cell cancer

Your diagnosis and treatment setting NCCP 00812b: metastatic or locally advanced cutaneous squamous cell carcinoma when curative surgery or curative radiotherapy is not appropriate. This guide covers skin SCC, not the separate cervical-cancer indication.

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Cemiplimab treatment

Active medicines: Cemiplimab

Emetogenic risk: Minimal

No routine prophylactic antiemetic order is populated for this phase. Agree an individual symptom-treatment plan where needed.

Additional phase-specific support

- Routine antiemetic prophylaxis or premedication is not usually required by the source. Provide an immune-toxicity contact/ alert plan; steroid or hormone treatment is prescribed only for the individual indication.
- Use effective contraception during cemiplimab treatment and for at least four months after the last dose. Discuss fertility before treatment, and follow the medicine-specific breastfeeding restrictions. Report a possible pregnancy promptly.
- Immunotherapy can cause inflammation in healthy organs during treatment or after it stops. Contact the oncology team promptly for new diarrhoea, blood in stools, cough or breathlessness, yellow skin or eyes, severe headache, vision changes, marked thirst, unusual weakness or a spreading rash. Tell any clinician that you are receiving or have received immunotherapy.
- This is not an issued prescription. The clinician and pharmacist must complete and authorise any supportive medicines.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00812 v3a · pSee current source monitoring/supportive-care sections · NCCP regimen and local supportive-care policy

Sources checked 2026-11-06 · SACTCheck v0.84.40 · Regimen-level drafts; prescriber and oncology-pharmacy verification required

Cemiplimab for advanced cutaneous squamous cell cancer

NCCP 00812 · Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Taking or receiving treatment

Cemiplimab is infused into a vein over 30 minutes in the source schedule. Allow additional time for assessment, preparation and observation. Tell staff immediately about itching, flushing, chills, pain, dizziness, breathlessness or swelling during the infusion.

STEP 2

Missed, vomited or held treatment

If a visit is missed or treatment is held, contact the unit for a new appointment and treatment decision. Do not assume the next scheduled date means treatment has been cleared to restart.

STEP 3

Appointments and supply

Arrange dates, medicine supply, transport and any needed interpreter or carer support. Ask whether bloods are taken in the oncology unit or elsewhere and how the result reaches the prescriber.

STEP 4

Paper copies and contact information

Keep this complete handbook, the three-page summary and the wallet card together. The QR code is optional. Ask the team to add your actual contacts and print a revised plan when treatment changes.

Sources: NCCP 00812 v3a: current regimen source. Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending.

Cemiplimab for advanced cutaneous squamous cell cancer

A record of what you experience

One cycle per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal CTCAE grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Cemiplimab is given by IV infusion on Day 1 every 21 days. The usual source dose is 350 mg over 30 minutes; the prescription and visit checks confirm your treatment. There is no fixed cycle total in this regimen: treatment continues while the team considers it beneficial and tolerable.

These pages are reusable recording sheets; the prescribed course length is stated above. Ask for additional sheets as needed.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

https://sactcheck.com/patient/00812/

Full patient information, sources and visual timeline. No identifiers are encoded.

Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

My diagnosis, plan and medicines

Complete these pages with your team. Leave unknown or inapplicable items blank. This record supports the consent discussion and your written prescription.

Diagnosis and overall plan

Original skin SCC site / pathology date

Stage / nodes / other sites / recurrence

Depth, nerve involvement or other high-risk features

Immunosuppression / transplant team if applicable

Earlier surgery / RT and reason curative local treatment is unsuitable

Medicines and safety

Anticancer medicines / actual dose / route / start date

Treatment received, missed or held / restart instruction

All other medicines / allergies / important previous reactions

Pregnancy-prevention or immune-alert plan / continuing precautions

Oncology / out-of-hours / pharmacy contacts

NCCN Guidelines for Patients: Squamous Cell Skin Cancer, 2023 - supplied education reference; original fields adapted to the Irish NCCP regimen context. No NCCN form or artwork reproduced.. Your Irish treating team confirms the diagnosis, prescription and follow-up plan.

My checks, symptoms and support

Record what each test is checking and when the result will be discussed. Your team explains which examinations or scans are needed for your diagnosis and treatment.

Test and purpose	Date / booking contact	Result discussion	Agreed next step

Record bloods, skin/lesion examinations, scans and specialist checks separately. Tumour-marker and scan intervals are individually selected; do not infer them from the cycle calendar.

Symptoms, skin and support

New symptoms / onset / change / action taken

Lesion or wound / pain / eating / weight / daily function

Skin self-check plan / optional dated photograph record kept privately

Sun protection / practical, emotional or family support needs

Questions, response review and the outcomes that matter to me

My urgent contact plan / what my carer should do

NCCN Guidelines for Patients: Squamous Cell Skin Cancer, 2023 - supplied education reference; original fields adapted to the Irish NCCP regimen context. No NCCN form or artwork reproduced.. Your Irish treating team confirms the diagnosis, prescription and follow-up plan.

Cemiplimab for advanced cutaneous squamous cell cancer

Cycle number: _____ Start date: _____ Next review: _____

Record the actual date, treatment received, any hold, important symptom and the next check. A review visit is not automatically a medicine dose or a completed cycle.

Cemiplimab: ongoing treatment visits · copy this sheet for another cycle of the same phase.

Cemiplimab · PD-1 checkpoint immunotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Bowels	☐ Usual pattern	☐ New change: report	☐ Persistent diarrhoea / constipation: call	☐ Blood, severe pain or dehydration: urgent review	☐ Collapse or severe sudden illness: 112/999
Breathing	☐ Usual breathing	☐ New cough / change: call	☐ Worsening breathlessness: urgent advice	☐ Breathless at rest: urgent assessment	☐ Severe breathing difficulty: 112/999
Skin	☐ No change	☐ New small change: report	☐ Persistent rash / itch: contact team	☐ Painful, spreading or blistering rash: urgent review	☐ Severe skin reaction with acute illness: emergency help
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Keep your treatment details to hand

NCCP 00812b: metastatic or locally advanced cutaneous squamous cell carcinoma when curative surgery or curative radiotherapy is not appropriate. This guide covers skin SCC, not the separate cervical-cancer indication. · NCCP 00812 v3a · Document revised 6 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Cemiplimab

Cemiplimab infusion every 3 weeks; no fixed cycle total in this regimen.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Use an ongoing dated visit record; there is no preset completion stamp total.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

SACTCheck v0.84.40 · NCCP 00812 v3a · Wallet-card sheet
Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.