Dabrafenib and trametinib for advanced BRAF-mutated melanoma

NCCP 00415 · Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Taking or receiving treatment

Swallow the prescribed capsules/tablet whole with water. Take each dose at least one hour before food or at least two hours after food. Take the once-daily trametinib dose at the same time each day, alongside either the morning or evening dabrafenib dose according to your written plan. Keep the prescribed storage instructions.

STEP 2

Missed, vomited or held treatment

The missed-dose windows differ: take missed dabrafenib only when more than six hours remain before the next dose; take missed trametinib only when more than 12 hours remain. Otherwise skip it. Do not double up. Do not take an extra trametinib dose after vomiting; contact the team or pharmacist about any vomited or uncertain dose.

STEP 3

Appointments and supply

Arrange dates, medicine supply, transport and any needed interpreter or carer support. Ask whether bloods are taken in the oncology unit or elsewhere and how the result reaches the prescriber.

STEP 4

Paper copies and contact information

Keep this complete handbook, the three-page summary and the wallet card together. The QR code is optional. Ask the team to add your actual contacts and print a revised plan when treatment changes.

Sources: NCCP 00415 v5: current regimen source. Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending.