SACTCHECK · NCCP 00563

My complete treatment handbook

Encorafenib and binimetinib for advanced BRAF-mutated melanoma

All information can be read on paper. Keep this with your treatment record.

> Use the urgent instructions and contact numbers in the following companion pages. Example telephone numbers must be replaced by your team before patient use. Your individual treatment plan takes precedence.

Contents

1. Your three-page treatment companion — page 2
2. Full treatment reference: medicines, schedule, tests, symptoms and follow-up — page 5
3. Detailed treatment and monitoring plan — page 26
4. Supportive medicines and care — page 30
5. Visits, access and practical arrangements — page 32
6. Your treatment passport and symptom record — page 33
7. Wallet card and phone instructions — page 37

SACTCheck v0.84.40 · Clinical, pharmacy and patient review pending. Formal consent is supplied separately.

Encorafenib and binimetinib for advanced BRAF-mutated melanoma

Daily targeted treatment for advanced BRAF V600-mutated melanoma.

NCCP 00563: adults with advanced unresectable or metastatic melanoma with a BRAF V600 mutation. · NCCP 00563 v4 · Document revised 6 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

NCCP 00563: adults with advanced unresectable or metastatic melanoma with a BRAF V600 mutation. This information supports individual consent; it does not issue or change a prescription.

My overall treatment

Encorafenib daily + binimetinib twice daily, continuously; reviewed regularly.

Continuous oral treatment
Continue while beneficial and tolerable, with team review; no fixed cycle total.

A 28-day reference calendar

Day 1–28
Every day in the reference calendar

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Baseline safety** NCCP / medicine-specific precautions	BRAF result; bloods, CK, BP, ECG, heart-pumping and skin checks.
During	**Bloods and organ checks** NCCP; local interval clarification where stated	Monthly bloods; CK/creatinine monthly for 6 months. ECG at 4 weeks then 12-weekly; heart check 3-monthly.
After / response	**Follow-up** Regimen / individual disease plan	Skin checks during treatment and for 6 months after; scans and further tests as planned.

Between treatments

- Both oral medicines continue daily unless the team tells you to pause.
- A review date does not automatically clear a held medicine to restart.

Bring the written medicine plan, test dates and actual oncology/out-of-hours contacts.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321
EXAMPLE ONLY — team must replace both numbers before patient use.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Energy / fever

Both medicines

Fatigue and headache can occur. Fever, marked weakness, dizziness or new neurological symptoms needs advice; do not assume it is a harmless drug effect.

3 Heart / circulation

Both medicines

Keep ECG and heart-pumping checks. Report palpitations, fainting, swollen ankles or new breathlessness; severe chest pain is an emergency.

5 Liver


Both medicines

Liver changes may have no early symptoms, so attend monthly tests. Report yellow skin or eyes, dark urine or upper-abdominal pain.

7 Muscles / strength

Binimetinib

CK can rise before or with muscle injury. Report muscle pain, weakness or dark urine, and keep the planned blood tests.



2 Eyes / vision

Both medicines

Blurred vision, spots, altered colour vision or eye pain needs prompt assessment. Retinal problems may occur with binimetinib.

4 Skin / new lesions

Both medicines

Report new growths or changes in existing spots and attend skin checks. Severe rash, blistering or peeling needs urgent help.

6 Kidneys / fluids

Binimetinib / combination

Kidney tests accompany CK monitoring. Reduced urine, dehydration or dark urine with muscle symptoms needs urgent advice.

8 Stomach / bowels

Both medicines

Nausea and diarrhoea may occur. Persistent vomiting, inability to drink, severe abdominal pain, black stools or bleeding needs assessment.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Contact the team promptly for new visual symptoms, chest pain, breathlessness, palpitations, faintness, severe muscle pain or weakness, dark urine, bleeding, severe rash, persistent diarrhoea or severe abdominal pain. Do not continue one medicine at an unchanged dose if the other has been held without a specific team instruction. Report markedly reduced urine, persistent vomiting, new muscle cramps or weakness, or palpitations promptly; the team may need to check salts, kidney function and tumour lysis risk.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Keep your treatment details to hand

NCCP 00563: adults with advanced unresectable or metastatic melanoma with a BRAF V600 mutation. · NCCP 00563 v4 · Document revised 6 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Encorafenib + Binimetinib

Encorafenib daily + binimetinib twice daily, continuously; reviewed regularly.

Phase: _____ Review plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

BACK / TREATMENT RECORD

My review record

Use a dated medicine/review record, including holds. Continuous treatment has no fixed completion stamp total.

Review	Date / initials

My treatment information

Latest review date: _____

Plan changed? Ask the team to update this card.

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each completed review. Daily tablets continue according to your prescribed plan; a review stamp does not mean the course is complete or authorise further treatment. Record interruptions or a changed plan. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

NCCP 00563 · VERSION 4

Your treatment, explained

NCCP 00563: adults with advanced unresectable or metastatic melanoma with a BRAF V600 mutation. · NCCP 00563 v4 · Document revised 6 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Encorafenib and binimetinib for advanced BRAF-mutated melanoma

Reference calendar for daily medicine and reviews

Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

ACT ON SYMPTOMS, NOT THE CALENDAR

When to get help

Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. Contact the team promptly for new visual symptoms, chest pain, breathlessness, palpitations, faintness, severe muscle pain or weakness, dark urine, bleeding, severe rash, persistent diarrhoea or severe abdominal pain. Do not continue one medicine at an unchanged dose if the other has been held without a specific team instruction. Report markedly reduced urine, persistent vomiting, new muscle cramps or weakness, or palpitations promptly; the team may need to check salts, kidney function and tumour lysis risk.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.

Your treatment and urgent-contact card is supplied separately and in the wallet-card section of the complete handbook.

Sources: nccp

YOUR COMPLETE REFERENCE

Read what you need, when you need it.

Start with the overview, explore the cycle, or go straight to a question. Urgent advice is always available.

How each drug works
The overall treatment plan
Your 28-day timeline
Baseline and cycle-specific checks
Whole treatment, scans and checks
Tablets, visits and practical planning
Understanding your diagnosis and treatment goal
Your medicines and the overall plan
Taking treatment and preparing for visits
Tests: what is required and what each result means
What to expect between visits
Skin checks, sun protection and optional photographs
Pregnancy, fertility and protecting others
Questions, decisions and practical support
Follow-up and your continuing record
Supportive medicines
Understanding the evidence
Questions to ask
Sources

UNDERSTAND YOUR TREATMENT


Medicines in this regimen

Encorafenib and binimetinib block two points in the BRAF-MEK growth pathway to help control eligible advanced melanoma. The team reviews response, symptoms and side effects to decide whether the treatment continues.

ORAL BRAF INHIBITOR

Encorafenib

Braftovi



Blocks altered BRAF signalling

Encorafenib blocks the BRAF V600 growth signal in eligible melanoma cells. Binimetinib acts further down the same pathway. They have different dose schedules and different safety decisions, so the prescription for one must not be assumed to apply to the other.

BRAF V600 confirmed → BRAF signal blocked → Partner MEK medicine

What to watch for


A change to binimetinib can change the encorafenib plan. Have interactions checked and avoid grapefruit.

Sources: nccp · nccn-patient

ORAL MEK INHIBITOR

Binimetinib

Mektovi



Blocks downstream MEK signalling

Binimetinib inhibits MEK in the cell-growth pathway. It can also affect normal muscle, heart and retinal tissues. CK blood tests, heart checks and prompt reporting of eye or muscle changes help the team identify problems before they become more serious.

MEK inhibited → Cancer-growth signal reduced → Muscle / heart / eye checks

What to watch for

Report vision change or muscle pain/weakness promptly; keep CK, kidney and heart checks even when you feel well.

Sources: nccp · nccn-patient

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

The anatomical side-effect map follows.

REGIMEN-SPECIFIC SIDE EFFECTS

Your anatomical side-effect map

A picture of side effects

Match each symptom box to its numbered body location. Tell your team about new symptoms.

1 Energy / fever

Both medicines

Fatigue and headache can occur. Fever, marked weakness, dizziness or new neurological symptoms needs advice; do not assume it is a harmless drug effect.

3 Heart / circulation

Both medicines

Keep ECG and heart-pumping checks. Report palpitations, fainting, swollen ankles or new breathlessness; severe chest pain is an emergency.

5 Liver


Both medicines

Liver changes may have no early symptoms, so attend monthly tests. Report yellow skin or eyes, dark urine or upper-abdominal pain.

7 Muscles / strength

Binimetinib

CK can rise before or with muscle injury. Report muscle pain, weakness or dark urine, and keep the planned blood tests.



2 Eyes / vision

Both medicines

Blurred vision, spots, altered colour vision or eye pain needs prompt assessment. Retinal problems may occur with binimetinib.

4 Skin / new lesions

Both medicines

Report new growths or changes in existing spots and attend skin checks. Severe rash, blistering or peeling needs urgent help.

6 Kidneys / fluids

Binimetinib / combination

Kidney tests accompany CK monitoring. Reduced urine, dehydration or dark urine with muscle symptoms needs urgent advice.

8 Stomach / bowels

Both medicines

Nausea and diarrhoea may occur. Persistent vomiting, inability to drink, severe abdominal pain, black stools or bleeding needs assessment.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

THE BIGGER TREATMENT PLAN

The overall treatment plan

DAILY MEDICINES

Continuous oral treatment

No preset cycle total **Encorafenib daily + binimetinib twice daily, continuously; reviewed regularly.**

Take encorafenib once daily and binimetinib twice daily, around 12 hours apart, continuously unless the team tells you to hold them. Treatment continues while beneficial and tolerable. A 28-day review calendar does not mean a routine break or a fixed total number of cycles.

The treatment setting determines duration; a reference calendar is not a prescription or a reason to restart a held medicine.

Sources: nccp

BASELINE → TREATMENT → FOLLOW-UP

Required safety checks and your individual disease-assessment plan

The team supplies actual dates and resolves any source ambiguity. A reference period does not replace a prescription, a local monitoring plan or an individual treatment decision.

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00563 v4: current regimen source · NCCN Guidelines for Patients: Melanoma, 2026 - supplied education reference

THE WHOLE TREATMENT JOURNEY

Encorafenib and binimetinib for advanced BRAF-mutated melanoma: treatment and checks

NCCP 00563: adults with advanced unresectable or metastatic melanoma with a BRAF V600 mutation.

Take encorafenib once daily and binimetinib twice daily, around 12 hours apart, continuously unless the team tells you to hold them. Treatment continues while beneficial and tolerable. A 28-day review calendar does not mean a routine break or a fixed total number of cycles.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	During treatment	Between visits	Response / follow-up	After a hold or stop
Bloods and organ safety NCCP 00563	Confirm BRAF V600 and treatment setting. Full blood count, kidney/liver tests, creatine kinase (CK) and creatinine, blood pressure, ECG, a heart-pumping assessment by echo or MUGA, and skin examination.	Blood count, kidney/liver tests monthly. CK and creatinine monthly for the first six months, then as indicated. ECG at four weeks, then every 12 weeks and after dose changes; heart-pumping assessment every three months. Skin assessment every 60 days, head/neck examination every three months and chest CT every six months. Response scans follow the individual plan.	Contact the team for new symptoms; extra tests may be needed between scheduled checks.	Safety bloods do not by themselves show whether cancer is responding.	Continue skin surveillance for six months after stopping as specified by the regimen. Agree continuing symptom/organ checks, the next melanoma assessment and medicine-specific pregnancy precautions.
Cancer, skin and daily function NCCP 00563	Record diagnosis, stage, current disease sites and the baseline assessment.	Keep the source-specific skin and clinical reviews and the individually agreed response assessments.	Record new lesions, lumps, wound changes, pain and effects on usual activities.	Write the next examination or scan date and the responsible team; no universal scan or tumour-marker interval is assumed.	Follow-up depends on stage, treatment and symptoms. Your team records the skin/lymph-node examination plan, whether body or brain imaging is needed, and which service owns each appointment. A general patient booklet does not set a universal scan interval for every patient.
Medicines and continuing precautions NCCP 00563	Review all medicines and the pregnancy/fertility plan.	Review treatment taken or received, holds, adverse effects and supportive medicines.	Keep the actual urgent contacts and do not restart a held medicine yourself.	Bring the medicine list and treatment summary to other services.	Use effective contraception during treatment and for at least one month after the last dose. Hormonal contraception needs an additional or alternative barrier method because interactions can reduce effectiveness. Discuss fertility beforehand and follow the team's breastfeeding instructions. Report a possible pregnancy promptly.

What each check is for

1

Continuous oral treatment

Daily medicines

Encorafenib and binimetinib block two points in the BRAF-MEK growth pathway to help control eligible advanced melanoma. The team reviews response, symptoms and side effects to decide whether the treatment continues.

Ask / act: Confirm the current treatment phase, prescription, tests and support medicines.

Sources: NCCP 00563

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-06 · Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.40

Day 1 → Day 28: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

Take encorafenib once daily and binimetinib twice daily, around 12 hours apart, continuously unless the team tells you to hold them. Treatment continues while beneficial and tolerable. A 28-day review calendar does not mean a routine break or a fixed total number of cycles.

Treatment timing: displayed days 1–28

Treatment or check	Days in this window	What happens
Daily oral treatment	Days 1–28	Encorafenib daily + binimetinib twice daily, continuously; reviewed regularly.
Safety reviews as scheduled	Days 1–28	Keep the specific blood, organ and skin-review appointments; they do not all follow the same interval.

Treatment or check	Days in this window	What happens
Symptoms and support	Days 1–28	Contact the team promptly for new visual symptoms, chest pain, breathlessness, palpitations, faintness, severe muscle pain or weakness, dark urine, bleeding, severe rash, persistent diarrhoea or severe abdominal pain. Do not continue one medicine at an unchanged dose if the other has been held without a specific team instruction. Report markedly reduced urine, persistent vomiting, new muscle cramps or weakness, or palpitations promptly; the team may need to check salts, kidney function and tumour lysis risk.

Timing note 1: Take encorafenib once daily and binimetinib twice daily, around 12 hours apart, continuously unless the team tells you to hold them. Treatment continues while beneficial and tolerable. A 28-day review calendar does not mean a routine break or a fixed total number of cycles.

Timing note 2: For a missed encorafenib dose, take it only if more than 12 hours remain before the next dose. For binimetinib, do not take a missed dose if fewer than six hours remain before the next; ask the pharmacist if timing is uncertain. Do not double doses. If either medicine is vomited, do not take an extra dose; take the next scheduled dose unless the team advises otherwise.

Timing note 3: Temporary holds may need coordinated changes to both medicines; no patient-led dose adjustment.

Timing note 4: The regimen is advanced melanoma; it is not an adjuvant treatment template.

Timing note 5: No trial outcome percentage is assigned without confirming the exact population, comparison and follow-up.

Timing note 6: Tumour lysis syndrome is a source-listed risk, especially with high tumour burden, kidney problems or dehydration. The team decides whether extra hydration and blood checks are needed.

> **Why these bands stay broad:** This calendar shows the pattern. Your team supplies actual dates, prescribed doses and any hold or restart instructions. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00563 v4: current regimen source ↗

> **There are no guaranteed "safe days".** The prescribed dose, treatment decision and local urgent-contact plan take priority. Call 112 or 999 for severe breathing difficulty, severe chest pain, collapse, a new seizure, sudden weakness or speech difficulty, or uncontrolled bleeding. Follow your oncology unit's written urgent-contact plan for other new or worsening symptoms; do not wait for the next appointment.

Complete timing and symptom notes

Day 1–28: Every day in the reference calendar

Encorafenib daily + binimetinib twice daily, continuously; reviewed regularly. Use the written medicine plan; report new symptoms and keep tests even during a treatment hold.

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Taking or receiving treatment

The usual method is to swallow the prescribed capsules/tablets whole with water, with or without food. If swallowing is difficult, ask the pharmacist for the source-permitted preparation: Braftovi capsules may be opened into about 20 mL of apple puree for immediate use, and Mektovi 15 mg tablets may be dispersed in about 10 mL of water, orange juice or apple juice, taken immediately, followed by a further 10 mL rinse. Do not assume this applies to other tablet strengths or improvise a preparation. Keep doses at consistent times. Avoid grapefruit and grapefruit juice, and have all medicines and supplements checked.

STEP 2

Missed, vomited or held treatment

For a missed encorafenib dose, take it only if more than 12 hours remain before the next dose. For binimetinib, do not take a missed dose if fewer than six hours remain before the next; ask the pharmacist if timing is uncertain. Do not double doses. If either medicine is vomited, do not take an extra dose; take the next scheduled dose unless the team advises otherwise.

STEP 3

Appointments and supply

Arrange dates, medicine supply, transport and any needed interpreter or carer support. Ask whether bloods are taken in the oncology unit or elsewhere and how the result reaches the prescriber.

STEP 4

Paper copies and contact information

Keep this complete handbook, the three-page summary and the wallet card together. The QR code is optional. Ask the team to add your actual contacts and print a revised plan when treatment changes.

Sources: NCCP 00563 v4: current regimen source. Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending.

MAKE THE PLAN PRACTICAL

Support around the medicines

1 An individual symptom plan

Ask which anti-sickness or other symptom medicines are prescribed, when to take them and when to call if they do not help. Do not start a borrowed medicine or alter treatment to manage a side effect yourself.

2 Your urgent contacts

Keep your actual oncology, out-of-hours and pharmacy contacts on the wallet card. Example numbers on a template must be replaced before you use it.

3 Pregnancy and fertility precautions

Use effective contraception during treatment and for at least one month after the last dose. Hormonal contraception needs an additional or alternative barrier method because interactions can reduce effectiveness. Discuss fertility beforehand and follow the team’s breastfeeding instructions. Report a possible pregnancy promptly.

4 Eating, activity and practical help

Report falling intake, weight loss, pain or fatigue early. The team can tailor nutrition, activity, wound, psychological and practical support to your needs.

5 Hydration and tumour lysis assessment

The team checks whether tumour burden, kidney problems or dehydration call for additional precautions and blood tests. Keep the individual fluid plan; do not force fluids if you have a restriction. Report reduced urine, persistent vomiting or marked weakness promptly.

Sources: nccp · nccn-patient

TREATMENT REFERENCE · 01

Understanding your diagnosis and treatment goal

Use these questions with your own pathology report and agreed treatment plan.

What the diagnosis says

Keep the pathology summary, original skin site, stage and details of any lymph-node involvement together. Breslow thickness and ulceration describe features of the original melanoma; they are different from the stage. If the diagnosis arose in the eye or a mucosal surface, ask whether this cutaneous-melanoma information applies.

Sources: nccn-patient

BRAF and other tumour tests

A BRAF V600 result helps decide whether the BRAF/MEK targeted medicines in this guide are suitable. This is usually a test of the tumour, not proof of an inherited family condition. Keep the actual result and test date; ask about genetic counselling only when your personal or family history makes that relevant.

Sources: nccn-patient

Surgery, radiotherapy and medicine have different roles

Record what surgery removed and whether the present medicine is postoperative treatment to lower recurrence risk or treatment of cancer that remains or has spread. Radiotherapy, stereotactic treatment or another operation may have a separate role; their timing and side effects need their own plan.

Sources: nccn-patient

Stage and response are different

Stage describes the extent of melanoma at diagnosis or a later reassessment. A response assessment asks how known cancer has changed with treatment. In the adjuvant setting there may be no visible tumour to measure: surveillance looks for recurrence while safety checks look for medicine effects.

Sources: nccn-patient

A new skin change still needs review

Continue to report a new or changing mole, a non-healing lesion, a new lump or an unexplained persistent symptom. Do not assume every new lesion is melanoma, and do not use a photograph or a good response at one site to rule out a problem elsewhere.

Sources: nccn-patient

TREATMENT REFERENCE · 02

Your medicines and the overall plan

Encorafenib and binimetinib block two points in the BRAF-MEK growth pathway to help control eligible advanced melanoma. The team reviews response, symptoms and side effects to decide whether the treatment continues.

The schedule in this setting

Take encorafenib once daily and binimetinib twice daily, around 12 hours apart, continuously unless the team tells you to hold them. Treatment continues while beneficial and tolerable. A 28-day review calendar does not mean a routine break or a fixed total number of cycles.

Sources: nccp

How the treatment works

Encorafenib blocks the BRAF V600 growth signal in eligible melanoma cells. Binimetinib acts further down the same pathway. They have different dose schedules and different safety decisions, so the prescription for one must not be assumed to apply to the other. Binimetinib inhibits MEK in the cell-growth pathway. It can also affect normal muscle, heart and retinal tissues. CK blood tests, heart checks and prompt reporting of eye or muscle changes help the team identify problems before they become more serious.

Sources: nccp

The role of earlier or later treatment

Record surgery, radiotherapy and other anticancer medicines separately, including dates and the specialist responsible. This schedule does not automatically include or authorise another treatment. Ask whether any overlap changes tests, symptoms to report or the timing of appointments.

Sources: nccn-patient

Pauses and stopping

A safety pause is an active treatment decision. Record which medicine is paused, the reason, the tests or recovery needed and who may authorise restarting. Do not restart from an old prescription or reset the history when a new phase begins.

Sources: nccp

TREATMENT REFERENCE · 03

Taking treatment and preparing for visits

Keep the written medicine plan beside this guide; it records your own dose and dates.

Taking or receiving the medicine

The usual method is to swallow the prescribed capsules/tablets whole with water, with or without food. If swallowing is difficult, ask the pharmacist for the source-permitted preparation: Braftovi capsules may be opened into about 20 mL of apple puree for immediate use, and Mektovi 15 mg tablets may be dispersed in about 10 mL of water, orange juice or apple juice, taken immediately, followed by a further 10 mL rinse. Do not assume this applies to other tablet strengths or improvise a preparation. Keep doses at consistent times. Avoid grapefruit and grapefruit juice, and have all medicines and supplements checked.

Sources: nccp

Missed, vomited or held treatment

For a missed encorafenib dose, take it only if more than 12 hours remain before the next dose. For binimetinib, do not take a missed dose if fewer than six hours remain before the next; ask the pharmacist if timing is uncertain. Do not double doses. If either medicine is vomited, do not take an extra dose; take the next scheduled dose unless the team advises otherwise.

Sources: nccp

Other medicines and supplements

Show the pharmacist a current list of prescriptions, over-the-counter medicines, vitamins, herbal products and supplements. Include anticoagulants, heart medicines, steroids and recent antibiotics. Check before adding, stopping or changing anything; an apparently harmless product can alter treatment or confuse a symptom assessment.

Sources: nccp

Practical preparation

Arrange medicine supply, transport, interpreter or carer support before it is needed. Bring the alert card, symptom record and questions. Ask which appointments require someone to accompany you and how to obtain urgent advice outside normal hours.

Sources: nccn-patient

TREATMENT REFERENCE · 04

Tests: what is required and what each result means

Safety tests, tumour tests and response scans answer different questions.

Before treatment

Confirm BRAF V600 and treatment setting. Full blood count, kidney/liver tests, creatine kinase (CK) and creatinine, blood pressure, ECG, a heart-pumping assessment by echo or MUGA, and skin examination.

Sources: nccp

During treatment

Blood count, kidney/liver tests monthly. CK and creatinine monthly for the first six months, then as indicated. ECG at four weeks, then every 12 weeks and after dose changes; heart-pumping assessment every three months. Skin assessment every 60 days, head/neck examination every three months and chest CT every six months. Response scans follow the individual plan.

Sources: nccp

Scans and tumour markers

Ask the team to write the next assessment and who will arrange it. Bloods used to monitor kidneys, liver, hormones or muscle are not automatically measures of cancer response. There is no universal tumour-marker blood schedule for these skin-cancer regimens; a scan or extra blood test is requested for a specific clinical reason.

Sources: nccp

After a pause or finish

Continue skin surveillance for six months after stopping as specified by the regimen. Agree continuing symptom/organ checks, the next melanoma assessment and medicine-specific pregnancy precautions.

Sources: nccp

Getting the result

For each test, record its purpose, due date, result and action. Agree how results reach you and who checks a result from another hospital. If an appointment or result has not arrived, contact the named service rather than assuming no news means no action is needed.

Sources: nccn-patient

TREATMENT REFERENCE · 05

What to expect between visits

You can feel well and still need the planned checks. New symptoms deserve attention even on a day without an appointment.

Your usual pattern

Record the start and change in symptoms, temperature if advised, bowel pattern, drinking, urine and the effect on normal tasks. A simple few-line record is enough. Use it to describe a change rather than trying to grade or diagnose a toxicity yourself.

Sources: nccp

When to call

Contact the team promptly for new visual symptoms, chest pain, breathlessness, palpitations, faintness, severe muscle pain or weakness, dark urine, bleeding, severe rash, persistent diarrhoea or severe abdominal pain. Do not continue one medicine at an unchanged dose if the other has been held without a specific team instruction. Report markedly reduced urine, persistent vomiting, new muscle cramps or weakness, or palpitations promptly; the team may need to check salts, kidney function and tumour lysis risk.

Sources: nccp

Eating, drinking and activity

Tell the team early about nausea, taste change, a sore mouth, diarrhoea or falling weight. Ask for an individual fluid and food plan if heart, kidney or other conditions limit intake. Gentle activity and planned rest can help maintain function; adjust to symptoms and ask for support when fatigue limits daily life.

Sources: nccn-patient

A held dose and a new phase

For a missed encorafenib dose, take it only if more than 12 hours remain before the next dose. For binimetinib, do not take a missed dose if fewer than six hours remain before the next; ask the pharmacist if timing is uncertain. Do not double doses. If either medicine is vomited, do not take an extra dose; take the next scheduled dose unless the team advises otherwise.

Sources: nccp

TREATMENT REFERENCE · 06

Skin checks, sun protection and optional photographs

This complements clinical assessment and does not replace it.

Protect your skin

Use shade, protective clothing, a hat and broad-spectrum sunscreen of at least SPF 30, and avoid tanning beds. Reapply sunscreen according to its instructions. Protection matters even when it is cloudy; choose a practical approach that lets you remain active safely.

Sources: nccn-patient

Know the pattern of your skin

Ask how to examine your skin, scars and relevant lymph-node areas and how often your own team recommends doing so. A mirror or a trusted helper can make hard-to-see areas easier to check. Report a new or changing lesion, a non-healing sore or a new lump rather than waiting for a routine visit.

Sources: nccn-patient

Photographs are optional

If useful, keep a dated photograph with consistent lighting and a size reference, following your team's advice. Photographs are optional and do not diagnose a cancer or replace examination. Keep them privately; this printable passport does not require you to upload images or other personal health information.

Sources: nccn-patient

Wounds, nails and sensitive skin

Use the dressings and skin-care plan recommended for your lesion and treatment. Ask before applying new creams or remedies to an open or treated area. Worsening pain, discharge, redness, bleeding or loss of function needs advice; routine skin-care instructions are not a substitute for assessment.

Sources: nccn-patient

TREATMENT REFERENCE · 07

Pregnancy, fertility and protecting others

Discuss these precautions before treatment even if pregnancy is not currently planned.

Medicine-specific precautions

Use effective contraception during treatment and for at least one month after the last dose. Hormonal contraception needs an additional or alternative barrier method because interactions can reduce effectiveness. Discuss fertility beforehand and follow the team's breastfeeding instructions. Report a possible pregnancy promptly.

Sources: nccp

Planning ahead

Ask early about fertility preservation, sexual health and symptoms affecting intimacy. Tell the team about any chance of pregnancy, breastfeeding or a partner's pregnancy so that the correct medicine-specific advice can be given. Keep the precautions and their end dates in the written plan.

Sources: nccp

Safe handling and supply

Do not share anticancer medicines. Keep oral medicines in their original labelled packaging, away from children and anyone who could take them accidentally. Return unused medicines to the supplying pharmacy and follow its handling advice. A treatment hold does not remove pregnancy precautions.

Sources: nccp

TREATMENT REFERENCE · 08

Questions, decisions and practical support

Your priorities and the burden of treatment belong in the discussion.

Benefits and uncertainties

Ask what evidence applies to your diagnosis, stage, biomarker and treatment setting, what outcome the team hopes to improve, and what may happen without this treatment. A trial average cannot predict an individual result; a benefit measured in one setting should not be transferred to another.

Sources: nccn-patient

Alternatives and review points

Discuss other suitable treatments, clinical trials if relevant, local treatment options and symptom-focused care. Ask when the decision will be reviewed and what would prompt a change. Supportive and palliative care can help with symptoms and quality of life alongside anticancer treatment.

Sources: nccn-patient

Help with daily life

Tell the team about work, transport, caring responsibilities, cost concerns, body image, anxiety or sleep. Ask which nurse, social worker, dietitian, rehabilitation professional or support organisation can help. Use resources relevant to your country and service; a US guide may describe different access or funding.

Sources: nccn-patient

Shared understanding

Ask someone on the team to check that you can explain the medicine schedule, next tests and urgent-contact route in your own words. An interpreter, large-print copy or a trusted supporter can help; none requires you to use a smartphone.

Sources: nccn-patient

TREATMENT REFERENCE · 09

Follow-up and your continuing record

Keep this record when treatment is paused, changed or completed.

Your follow-up plan

Follow-up depends on stage, treatment and symptoms. Your team records the skin/lymph-node examination plan, whether body or brain imaging is needed, and which service owns each appointment. A general patient booklet does not set a universal scan interval for every patient.

Sources: nccn-patient

Continuing medicine precautions

Continue skin surveillance for six months after stopping as specified by the regimen. Agree continuing symptom/organ checks, the next melanoma assessment and medicine-specific pregnancy precautions.

Sources: nccp

Treatment summary for other clinicians

Record the medicines and last dates, important reactions, any long-term hormone replacement or steroid taper, and the responsible oncology contact. Show this information in urgent care and to other specialists. A different treatment phase does not erase earlier toxicity.

Sources: nccp

The complete paper version

This handbook can be used without a phone: keep the schedule, symptom advice, medicine list, tests and contacts together. The QR code is an optional route to the matching regimen information. Ask the team to print updates when your plan changes and keep the latest dated copy.

Sources: nccn-patient

What the evidence means for your treatment

The evidence discussion must match the diagnosis, biomarker and treatment setting shown in this guide.

Ask which outcomes matter in your setting and what benefit is realistic for you. No survival percentage from another regimen, dose combination or disease setting is assigned here.

Sources: nccp · nccn-patient

PREPARE FOR THE CONVERSATION

Questions worth asking

What is the aim of my treatment?

Encorafenib and binimetinib block two points in the BRAF-MEK growth pathway to help control eligible advanced melanoma. The team reviews response, symptoms and side effects to decide whether the treatment continues.

Is there a fixed number of cycles?

Take encorafenib once daily and binimetinib twice daily, around 12 hours apart, continuously unless the team tells you to hold them. Treatment continues while beneficial and tolerable. A 28-day review calendar does not mean a routine break or a fixed total number of cycles.

What happens if treatment is held?

For a missed encorafenib dose, take it only if more than 12 hours remain before the next dose. For binimetinib, do not take a missed dose if fewer than six hours remain before the next; ask the pharmacist if timing is uncertain. Do not double doses. If either medicine is vomited, do not take an extra dose; take the next scheduled dose unless the team advises otherwise.

When are my next tests?

Blood count, kidney/liver tests monthly. CK and creatinine monthly for the first six months, then as indicated. ECG at four weeks, then every 12 weeks and after dose changes; heart-pumping assessment every three months. Skin assessment every 60 days, head/neck examination every three months and chest CT every six months. Response scans follow the individual plan. Your team adds the actual dates.

Can I use this without a smartphone?

Yes. The full handbook, three-page summary and wallet card are printable. The QR code is optional; keep actual contacts and written appointments on the paper copy.

> **Try explaining the plan back:** Can you explain your treatment setting, medicine schedule, next tests, the symptoms that need urgent advice and the number you would call?

SOURCE TRANSPARENCY

Where this information comes from

Regimen schedule: NCCP 00563 v4. Medicine explanations and safety information: sources below. The timeline is an educational illustration, not a measured risk curve or an individual prediction.

- NCCP 00563 v4: current regimen source
- NCCN Guidelines for Patients: Melanoma, 2026 - supplied education reference

SACTCheck · Developed by Dr Paul O'Brien · Sources checked 2026-11-06.
This information supports your treatment discussion. Your oncology team confirms your personal treatment and urgent-care plan.

Written source addresses

https://healthservice.hse.ie/documents/6744/563_Encorafenib-and-Binimetinib-Therapy.pdf

https://www.nccn.org/patients/guidelines/content/PDF/melanoma-patient.pdf

BASELINE → TREATMENT → FOLLOW-UP

Required safety checks and your individual disease-assessment plan

The team supplies actual dates and resolves any source ambiguity. A reference period does not replace a prescription, a local monitoring plan or an individual treatment decision.

Check	Before starting	During treatment	Between visits	Response / follow-up	After a hold or stop
Bloods and organ safety	Confirm BRAF V600 and treatment setting. Full blood count, kidney/liver tests, creatine kinase (CK) and creatinine, blood pressure, ECG, a heart-pumping assessment by echo or MUGA, and skin examination.	Blood count, kidney/liver tests monthly. CK and creatinine monthly for the first six months, then as indicated. ECG at four weeks, then every 12 weeks and after dose changes; heart-pumping assessment every three months. Skin assessment every 60 days, head/neck examination every three months and chest CT every six months. Response scans follow the individual plan.	Contact the team for new symptoms; extra tests may be needed between scheduled checks.	Safety bloods do not by themselves show whether cancer is responding.	Continue skin surveillance for six months after stopping as specified by the regimen. Agree continuing symptom/organ checks, the next melanoma assessment and medicine-specific pregnancy precautions.
Cancer, skin and daily function	Record diagnosis, stage, current disease sites and the baseline assessment.	Keep the source-specific skin and clinical reviews and the individually agreed response assessments.	Record new lesions, lumps, wound changes, pain and effects on usual activities.	Write the next examination or scan date and the responsible team; no universal scan or tumour-marker interval is assumed.	Follow-up depends on stage, treatment and symptoms. Your team records the skin/lymph-node examination plan, whether body or brain imaging is needed, and which service owns each appointment. A general patient booklet does not set a universal scan interval for every patient.
Medicines and continuing precautions	Review all medicines and the pregnancy/fertility plan.	Review treatment taken or received, holds, adverse effects and supportive medicines.	Keep the actual urgent contacts and do not restart a held medicine yourself.	Bring the medicine list and treatment summary to other services.	Use effective contraception during treatment and for at least one month after the last dose. Hormonal contraception needs an additional or alternative barrier method because interactions can reduce effectiveness. Discuss fertility beforehand and follow the team's breastfeeding instructions. Report a possible pregnancy promptly.

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00563 v4: current regimen source · NCCN Guidelines for Patients: Melanoma, 2026 - supplied education reference

THE WHOLE TREATMENT JOURNEY

Encorafenib and binimetinib for advanced BRAF-mutated melanoma: treatment and checks

NCCP 00563: adults with advanced unresectable or metastatic melanoma with a BRAF V600 mutation.

Take encorafenib once daily and binimetinib twice daily, around 12 hours apart, continuously unless the team tells you to hold them. Treatment continues while beneficial and tolerable. A 28-day review calendar does not mean a routine break or a fixed total number of cycles.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	During treatment	Between visits	Response / follow-up	After a hold or stop
Bloods and organ safety NCCP 00563	Confirm BRAF V600 and treatment setting. Full blood count, kidney/liver tests, creatine kinase (CK) and creatinine, blood pressure, ECG, a heart-pumping assessment by echo or MUGA, and skin examination.	Blood count, kidney/liver tests monthly. CK and creatinine monthly for the first six months, then as indicated. ECG at four weeks, then every 12 weeks and after dose changes; heart-pumping assessment every three months. Skin assessment every 60 days, head/neck examination every three months and chest CT every six months. Response scans follow the individual plan.	Contact the team for new symptoms; extra tests may be needed between scheduled checks.	Safety bloods do not by themselves show whether cancer is responding.	Continue skin surveillance for six months after stopping as specified by the regimen. Agree continuing symptom/organ checks, the next melanoma assessment and medicine-specific pregnancy precautions.
Cancer, skin and daily function NCCP 00563	Record diagnosis, stage, current disease sites and the baseline assessment.	Keep the source-specific skin and clinical reviews and the individually agreed response assessments.	Record new lesions, lumps, wound changes, pain and effects on usual activities.	Write the next examination or scan date and the responsible team; no universal scan or tumour-marker interval is assumed.	Follow-up depends on stage, treatment and symptoms. Your team records the skin/lymph-node examination plan, whether body or brain imaging is needed, and which service owns each appointment. A general patient booklet does not set a universal scan interval for every patient.

Care / check	Before starting	During treatment	Between visits	Response / follow-up	After a hold or stop
Medicines and continuing precautions NCCP 00563	Review all medicines and the pregnancy/fertility plan.	Review treatment taken or received, holds, adverse effects and supportive medicines.	Keep the actual urgent contacts and do not restart a held medicine yourself.	Bring the medicine list and treatment summary to other services.	Use effective contraception during treatment and for at least one month after the last dose. Hormonal contraception needs an additional or alternative barrier method because interactions can reduce effectiveness. Discuss fertility beforehand and follow the team's breastfeeding instructions. Report a possible pregnancy promptly.

What each check is for

1

Continuous oral treatment

Daily medicines

Encorafenib and binimetinib block two points in the BRAF-MEK growth pathway to help control eligible advanced melanoma. The team reviews response, symptoms and side effects to decide whether the treatment continues.

Ask / act: Confirm the current treatment phase, prescription, tests and support medicines.

Sources: NCCP 00563

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-06 · Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.40

Encorafenib and binimetinib for advanced BRAF-mutated melanoma

Your diagnosis and treatment setting NCCP 00563: adults with advanced unresectable or metastatic melanoma with a BRAF V600 mutation.

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Separate scripts for each phase

Choose the active phase in the clinician workflow. Each script has its own medicines, timing and review checks.

- **Continuous oral treatment** — Moderate to high (oral)
  Encorafenib + Binimetinib

https://sactcheck.com/patient/00563/supportive-care.html

Encorafenib and binimetinib for advanced BRAF-mutated melanoma

Your diagnosis and treatment setting NCCP 00563: adults with advanced unresectable or metastatic melanoma with a BRAF V600 mutation.

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Continuous oral treatment

Active medicines: Encorafenib + Binimetinib

Emetogenic risk: Moderate to high (oral)

Medicine	Dose / route	When
Individual anti-sickness / rescue plan	To be completed by the prescribing team if needed	As specified in the individual plan; no routine schedule issued here Source emetogenic category and local policy guide the decision. No medicine, dose or administration order is selected by this draft.

Additional phase-specific support

- Assess tumour lysis risk when tumour burden, kidney problems or dehydration raise concern; fluids and additional tests are individual prescriptions. Both oral components are moderate-to-high emetic risk in the source.
- Use effective contraception during treatment and for at least one month after the last dose. Hormonal contraception needs an additional or alternative barrier method because interactions can reduce effectiveness. Discuss fertility beforehand and follow the team's breastfeeding instructions. Report a possible pregnancy promptly.
- Contact the team promptly for new visual symptoms, chest pain, breathlessness, palpitations, faintness, severe muscle pain or weakness, dark urine, bleeding, severe rash, persistent diarrhoea or severe abdominal pain. Do not continue one medicine at an unchanged dose if the other has been held without a specific team instruction. Report markedly reduced urine, persistent vomiting, new muscle cramps or weakness, or palpitations promptly; the team may need to check salts, kidney function and tumour lysis risk.
- This is not an issued prescription. The clinician and pharmacist must complete and authorise any supportive medicines.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00563 v4 · p8–10 · NCCP regimen and local supportive-care policy

Sources checked 2026-11-06 · SACTCheck v0.84.40 · Regimen-level drafts; prescriber and oncology-pharmacy verification required

Encorafenib and binimetinib for advanced BRAF-mutated melanoma

NCCP 00563 · Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Taking or receiving treatment

The usual method is to swallow the prescribed capsules/tablets whole with water, with or without food. If swallowing is difficult, ask the pharmacist for the source-permitted preparation: Braftovi capsules may be opened into about 20 mL of apple puree for immediate use, and Mektovi 15 mg tablets may be dispersed in about 10 mL of water, orange juice or apple juice, taken immediately, followed by a further 10 mL rinse. Do not assume this applies to other tablet strengths or improvise a preparation. Keep doses at consistent times. Avoid grapefruit and grapefruit juice, and have all medicines and supplements checked.

STEP 2

Missed, vomited or held treatment

For a missed encorafenib dose, take it only if more than 12 hours remain before the next dose. For binimetinib, do not take a missed dose if fewer than six hours remain before the next; ask the pharmacist if timing is uncertain. Do not double doses. If either medicine is vomited, do not take an extra dose; take the next scheduled dose unless the team advises otherwise.

STEP 3

Appointments and supply

Arrange dates, medicine supply, transport and any needed interpreter or carer support. Ask whether bloods are taken in the oncology unit or elsewhere and how the result reaches the prescriber.

STEP 4

Paper copies and contact information

Keep this complete handbook, the three-page summary and the wallet card together. The QR code is optional. Ask the team to add your actual contacts and print a revised plan when treatment changes.

Sources: NCCP 00563 v4: current regimen source. Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending.

Encorafenib and binimetinib for advanced BRAF-mutated melanoma

A record of what you experience

One review period per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal CTCAE grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

Take encorafenib once daily and binimetinib twice daily, around 12 hours apart, continuously unless the team tells you to hold them. Treatment continues while beneficial and tolerable. A 28-day review calendar does not mean a routine break or a fixed total number of cycles.

These pages are reusable recording sheets; the prescribed course length is stated above. Ask for additional sheets as needed.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital's instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

https://sactcheck.com/patient/00563/

Full patient information, sources and visual timeline. No identifiers are encoded.

Draft source reconciliation; independent clinical, pharmacy and patient/PPI review pending

My diagnosis, plan and medicines

Complete these pages with your team. Leave unknown or inapplicable items blank. This record supports the consent discussion and your written prescription.

Diagnosis and overall plan

Original melanoma site / pathology date

Stage / Breslow thickness / ulceration

Lymph-node or other disease sites

BRAF result / date / other relevant tumour tests

Advanced or adjuvant intent / earlier surgery or RT

Medicines and safety

Anticancer medicines / actual dose / route / start date

Treatment received, missed or held / restart instruction

All other medicines / allergies / important previous reactions

Pregnancy-prevention or immune-alert plan / continuing precautions

Oncology / out-of-hours / pharmacy contacts

NCCN Guidelines for Patients: Melanoma, 2026 - supplied education reference; original fields adapted to the Irish NCCP regimen context. No NCCN form or artwork reproduced.. Your Irish treating team confirms the diagnosis, prescription and follow-up plan.

My checks, symptoms and support

Record what each test is checking and when the result will be discussed. Your team explains which examinations or scans are needed for your diagnosis and treatment.

Test and purpose	Date / booking contact	Result discussion	Agreed next step

Record bloods, skin/lesion examinations, scans and specialist checks separately. Tumour-marker and scan intervals are individually selected; do not infer them from the cycle calendar.

Symptoms, skin and support

New symptoms / onset / change / action taken

Lesion or wound / pain / eating / weight / daily function

Skin self-check plan / optional dated photograph record kept privately

Sun protection / practical, emotional or family support needs

Questions, response review and the outcomes that matter to me

My urgent contact plan / what my carer should do

NCCN Guidelines for Patients: Melanoma, 2026 - supplied education reference; original fields adapted to the Irish NCCP regimen context. No NCCN form or artwork reproduced.. Your Irish treating team confirms the diagnosis, prescription and follow-up plan.

Encorafenib and binimetinib for advanced BRAF-mutated melanoma

Review period: _____ Start date: _____ Next review: _____

Record the actual date, treatment received, any hold, important symptom and the next check. A review visit is not automatically a medicine dose or a completed cycle.

Daily medicines and review record · copy this sheet for another review period of the same phase.

Encorafenib · Oral BRAF inhibitor

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Skin	☐ No change	☐ New small change: report	☐ Persistent rash / itch: contact team	☐ Painful, spreading or blistering rash: urgent review	☐ Severe skin reaction with acute illness: emergency help
Nausea	☐ None	☐ Queasy but eating and drinking	☐ Intake reduced: contact the team	☐ Unable to maintain intake: urgent review	☐ Collapse / severe dehydration: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Binimetinib · Oral MEK inhibitor

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Eyes	☐ Usual vision	☐ New eye symptom: contact team	☐ Persistent eye pain / blur: urgent advice	☐ Sudden vision loss: immediate urgent assessment	☐ Vision change with stroke signs: 112/999
Muscles	☐ Usual strength	☐ New aches: report	☐ Pain / weakness limiting tasks: call	☐ Marked weakness or dark urine: urgent review	☐ Breathing / swallowing weakness or collapse: 112/999
Breathing	☐ Usual breathing	☐ New cough / change: call	☐ Worsening breathlessness: urgent advice	☐ Breathless at rest: urgent assessment	☐ Severe breathing difficulty: 112/999
Bowels	☐ Usual pattern	☐ New change: report	☐ Persistent diarrhoea / constipation: call	☐ Blood, severe pain or dehydration: urgent review	☐ Collapse or severe sudden illness: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

Keep your treatment details to hand

NCCP 00563: adults with advanced unresectable or metastatic melanoma with a BRAF V600 mutation. · NCCP 00563 v4 · Document revised 6 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Encorafenib + Binimetinib

Encorafenib daily + binimetinib twice daily, continuously; reviewed regularly.

Phase: _____ Review plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

BACK / TREATMENT RECORD

My review record

Use a dated medicine/review record, including holds. Continuous treatment has no fixed completion stamp total.

Review	Date / initials

My treatment information

Latest review date: _____

Plan changed? Ask the team to update this card.

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each completed review. Daily tablets continue according to your prescribed plan; a review stamp does not mean the course is complete or authorise further treatment. Record interruptions or a changed plan. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

SACTCheck v0.84.40 · NCCP 00563 v4 · Wallet-card sheet
Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.