# Cisplatin + etoposide with radiotherapy

Chemotherapy and radiation work together to treat limited-stage small-cell lung cancer. Your team explains your treatment aim and choices.

> For limited-stage lung cancer: IV cisplatin/etoposide every 21 days. Oral, split-dose, carboplatin or 28-day plans need personalised information.

## My overall treatment

Reference IV plan: four 21-day cycles. Day 1: cisplatin + etoposide. Days 2–3: etoposide. Radiation follows its own appointments. Capsules, split cisplatin, carboplatin substitution or 28-day cycles require a different personalised prescription.

**Chemotherapy**
Three treatment days per reference cycle, then recovery. Your team confirms each cycle after safety checks.

**Radiotherapy**
Usually starts with cycle 1. Radiation oncology sets the dose and visits; chemotherapy delays need separate coordination.

**After treatment**
Your team reviews recovery, persistent symptoms and response. Scan or marker timing is individual.

## A typical 21-day cycle

| Day 1 | Day 2–3 | Day 4–7 | Day 8–21 |
|---|---|---|---|
| reference combination | Days 2–3: etoposide | Support and recovery | Between chemotherapy cycles |

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

## My tests and checks

| When | What is checked | Timing / reason |
|---|---|---|
| **Before** | **Blood count, kidneys and liver**<br>Required by regimen | Before treatment |
| **Before** | **Audiology / creatinine clearance**<br>If clinically indicated | When clinically indicated |
| **During** | **Bloods, hearing and nerves**<br>Required by regimen | Before every chemotherapy cycle |
| **After / review** | **Recovery and response**<br>Team-agreed plan | Team-agreed timing; no universal interval |

## Between treatments

- Follow the prescribed anti-sickness course and fluid advice; call early if eating or drinking becomes difficult.
- Fever, new breathlessness, uncontrolled vomiting or inability to drink needs urgent advice, including after treatment ends.

Confirm your prescription, radiation appointments and blood tests. Bring your medicine list and daytime / overnight contact numbers.

**When to get help**

**Call 112 / 999** for severe breathing difficulty, severe chest pain, collapse or a seizure.

**Call oncology immediately:** temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment.

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE NUMBERS ONLY — your team must replace these before patient use.**


Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

# What to watch for

You do not need to know which medicine caused a symptom before calling.

### 1. Hearing / nerves

Cisplatin

Report ringing, reduced hearing, numbness or weakness before the next dose.

### 3. Breathing


Chemoradiation

New or worsening breathlessness needs urgent assessment, including after treatment finishes.

### 5. Kidneys / fluid balance

Cisplatin

Blood tests and fluids protect kidneys. Report much less urine or poor intake.




### 2. Mouth / swallowing

Treatment and chest radiation

Soreness may affect food and fluids. Contact the team early; inability to drink is urgent.

### 4. Sickness / appetite

Chemotherapy

Follow the anti-sickness plan. Call if vomiting continues or fluids will not stay down.

### 6. Blood cells / IV site

Chemotherapy

Fever or sudden illness can be serious. Report bleeding or infusion pain promptly.

Selected effects, not every possible effect or a prediction. Match each number to its symptom box. Blood, skin and nerve markers are schematic.

Call oncology immediately for fever, shaking chills, sudden illness, new breathlessness, uncontrolled vomiting, inability to swallow fluids, much less urine or unusual bleeding. Severe breathing difficulty, chest pain, collapse or heavy bleeding needs emergency care. Tell staff immediately about infusion pain, swelling or allergy symptoms.

**When to get help**

**Call 112 / 999** for severe breathing difficulty, severe chest pain, collapse or a seizure.

**Call oncology immediately:** temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment.

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE NUMBERS ONLY — your team must replace these before patient use.**


Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

# Keep your treatment details to hand

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

**FRONT / CONTACTS**

## My Treatment Card

SACTCheck

Name: _____

**Cisplatin + etoposide / RT**

Day 1: both drugs. Days 2–3: etoposide. Usually every 21 days ×4; RT plan separate.

Phase: _____ Planned cycles: 4

**Daytime (EXAMPLE)** 087 12345678

**Out of hours (EXAMPLE)** 087 87654321

**Unwell or fever? Call oncology now.**
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

**BACK / TREATMENT RECORD**

## My cycle record

Team stamps after each completed cycle.

C1 Date / initials

C2 Date / initials

C3 Date / initials

C4 Date / initials

My treatment information

Plan changed? Ask the team to update this card.

**FOLD centre — do not cut**

## Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

## Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

### iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

### Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

## Your local contacts and resources

**Oncology daytime:** 087 12345678 · **Out of hours:** 087 87654321

**EXAMPLE NUMBERS ONLY — your team must replace these before patient use.**

Hospital / team: _____ Backup contact route: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.


Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.