

Modified FOLFOX-6 (mFOLFOX6)

Colorectal / gastrointestinal cancer

Where changes may appear

The body systems and labels below are a memory aid. They are regimen/agent-specific and do not predict which effects you will experience.

Oxaliplatin

Peripheral neuropathy / cold sensitivity	Tingling, numbness or pain, often triggered by cold. Avoid cold exposure if advised and report symptoms before the next dose.
Acute throat or jaw sensation	A tight throat, jaw discomfort or difficulty taking a breath can occur soon after oxaliplatin. Tell staff immediately; severe breathing difficulty is an emergency.
Hypersensitivity reaction	Flushing, rash, wheeze, swelling or dizziness during/after infusion needs immediate assessment.

5-FU + leucovorin

Diarrhoea and mouth inflammation	Persistent diarrhoea, mouth ulcers or inability to drink can lead to dehydration and needs prompt oncology advice.
Hand-foot skin reaction	Soreness, redness, peeling or pain of palms/soles should be reported.
Rare heart toxicity	New chest pain, pressure, palpitations or breathlessness during 5-FU needs urgent assessment.

Regimen-wide

Low blood counts / infection	Fever, shivering or sudden illness can occur when white cells are low. Follow your oncology SOS temperature instructions.
Severe fluoropyrimidine toxicity	Unexpectedly severe diarrhoea, mouth ulcers, confusion or rapid deterioration can signal serious 5-FU toxicity and needs urgent review.

When to call: follow your oncology SOS instructions for fever, rapidly worsening symptoms, new breathing difficulty, persistent diarrhoea, significant bleeding, severe rash, jaundice, confusion or other concerning changes. For a medical emergency in Ireland call 112/999.



Scan for this regimen portal: <https://sactcheck.com/docs/patient/00209/>

Clinical-review prototype. Your current oncology plan and local emergency instructions take priority.