

# Pembrolizumab + pemetrexed + carboplatin

Metastatic non-squamous NSCLC

## Where changes may appear

The body systems and labels below are a memory aid. They are regimen/agent-specific and do not predict which effects you will experience.

## Pembrolizumab - immune effects

|                                                   |                                                                                                                                     |
|---------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------|
| <b>Lung inflammation (pneumonitis)</b>            | New cough or breathlessness needs prompt assessment.                                                                                |
| <b>Bowel / liver inflammation</b>                 | Persistent diarrhoea, abdominal pain, yellow eyes or dark urine can signal immune inflammation.                                     |
| <b>Hormone or sugar changes</b>                   | Marked fatigue, dizziness, headache, thirst or passing much more urine can reflect thyroid, adrenal, pituitary or glucose problems. |
| <b>Skin, kidney, nerve or muscle inflammation</b> | Spreading rash, blistering, reduced urine, new weakness, numbness or visual change needs assessment.                                |

## Pemetrexed

|                                     |                                                                                                                       |
|-------------------------------------|-----------------------------------------------------------------------------------------------------------------------|
| <b>Kidney toxicity</b>              | Kidney function is checked because pemetrexed clearance depends on it. Report reduced urine, swelling or dehydration. |
| <b>Mouth / gut and skin effects</b> | Mouth soreness, diarrhoea or rash can occur; folic acid and vitamin B12 support are important.                        |

## Carboplatin + chemotherapy

|                                            |                                                                                                            |
|--------------------------------------------|------------------------------------------------------------------------------------------------------------|
| <b>Low blood counts / infection</b>        | Fever, shivering, bleeding or unusual breathlessness can reflect low blood counts and needs prompt advice. |
| <b>Nausea, fatigue and appetite change</b> | Use prescribed supportive medicines and contact the team if you cannot maintain fluids.                    |
| <b>Hypersensitivity reaction</b>           | Rash, wheeze, swelling or dizziness during infusion needs immediate assessment.                            |

**When to call:** follow your oncology SOS instructions for fever, rapidly worsening symptoms, new breathing difficulty, persistent diarrhoea, significant bleeding, severe rash, jaundice, confusion or other concerning changes. For a medical emergency in Ireland call 112/999.



Scan for this regimen portal: <https://sactcheck.com/docs/patient/00568/>

Clinical-review prototype. Your current oncology plan and local emergency instructions take priority.