

Adjuvant abemaciclib + endocrine therapy

High-risk HR-positive / HER2-negative early breast cancer

Where changes may appear

The body systems and labels below are a memory aid. They are regimen/agent-specific and do not predict which effects you will experience.

Abemaciclib

Diarrhoea	Can begin early. Follow the team's diarrhoea plan, maintain fluids and contact oncology for persistent/severe symptoms or dehydration.
Low neutrophils / infection	Fever, shivering or sudden illness needs urgent oncology contact.
Venous blood clot (VTE)	One-sided leg swelling/pain, sudden breathlessness or chest pain needs urgent assessment.
Lung inflammation (ILD/pneumonitis)	New or worsening cough or breathlessness needs prompt assessment.
Liver toxicity	Liver blood tests are monitored; jaundice, dark urine or marked worsening fatigue should be reported.

Endocrine therapy

Agent-specific effects	The endocrine medicine may be an aromatase inhibitor or tamoxifen depending on the plan. Hot flushes, joint symptoms, vaginal symptoms and bone-health issues vary by agent.
Do not assume all risks are identical	Ask which endocrine agent you are taking and use its medicine-specific information alongside this portal.

When to call: follow your oncology SOS instructions for fever, rapidly worsening symptoms, new breathing difficulty, persistent diarrhoea, significant bleeding, severe rash, jaundice, confusion or other concerning changes. For a medical emergency in Ireland call 112/999.



Scan for this regimen portal: <https://sactcheck.com/docs/patient/00619/>

Clinical-review prototype. Your current oncology plan and local emergency instructions take priority.