

Pembrolizumab + carboplatin/paclitaxel -> AC

Early triple-negative breast cancer

Where changes may appear

The body systems and labels below are a memory aid. They are regimen/agent-specific and do not predict which effects you will experience.

Pembrolizumab - immune effects

Pneumonitis	New cough or breathlessness needs prompt assessment.
Colitis / hepatitis	Persistent diarrhoea, abdominal pain, jaundice or dark urine can signal immune inflammation.
Endocrine toxicity	Marked fatigue, dizziness, headache, thirst or excess urination can reflect hormone or glucose disturbance.
Skin, kidney, nerve or muscle inflammation	Spreading rash, blistering, reduced urine, weakness, numbness or visual change needs assessment.

Paclitaxel + carboplatin phase

Peripheral neuropathy	New tingling, numbness, burning pain or difficulty with fine tasks should be reported.
Hypersensitivity / infusion reaction	Flushing, rash, wheeze, swelling or dizziness during infusion needs immediate assessment.
Low blood counts / infection	Fever, shivering, bleeding or sudden illness needs prompt oncology contact.

Doxorubicin + cyclophosphamide phase

Heart toxicity	New chest pain, breathlessness, swelling or palpitations needs assessment.
Urine / bladder changes	Doxorubicin can temporarily redden urine; painful urination or visible blood should still be reported.
Mouth, nausea and blood-count effects	Mouth ulcers, vomiting despite medicines, fever or bleeding should be reported promptly.

When to call: follow your oncology SOS instructions for fever, rapidly worsening symptoms, new breathing difficulty, persistent diarrhoea, significant bleeding, severe rash, jaundice, confusion or other concerning changes. For a medical emergency in Ireland call 112/999.



Scan for this regimen portal: <https://sactcheck.com/docs/patient/00857/>

Clinical-review prototype. Your current oncology plan and local emergency instructions take priority.