Triptorelin 11.25 mg: hormone injection every 12 weeks

National Cancer Control Programme (NCCP) 00480 · Source reconciled draft; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Before starting

Full blood count, kidney and liver tests, blood glucose and a bone blood profile. Review heart disease, blood pressure, diabetes, mood, falls and fracture risk, sexual health, medicines and the treatment goal.

STEP 2

Taking or receiving treatment

A trained healthcare professional prepares and gives the injection using the product-specific instructions. Confirm the 11.25 mg formulation, agreed route and next appointment. Other triptorelin strengths have different schedules.

STEP 3

Tests and reviews

Repeat blood count, kidney and liver tests, glucose and bone blood profile when clinically indicated. Review symptoms, PSA and testosterone; scans and the frequency of tests follow the individual plan.

STEP 4

Specific instructions

The team assesses temporary anti-androgen protection against the initial hormone flare. Report worsening pain or trouble passing urine promptly. New weakness or changes in bladder/bowel control require emergency assessment. Keep the written injection dates even if you feel well.

STEP 5

Leaving with a clear plan

Confirm the exact next dose or appointment, duration and verified daytime and out-of-hours contacts.

Sources: NCCP 00480 v3: official regimen. Source reconciled draft; independent clinical, pharmacy and patient/PPI review pending.