Relugolix: daily hormone treatment

National Cancer Control Programme (NCCP) 00830 · Source reconciled draft; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Before starting

Full blood count, kidney and liver tests, blood glucose and a bone blood profile. Review heart disease, blood pressure, diabetes, mood, falls and fracture risk, sexual health, medicines and the treatment goal. An electrocardiogram (ECG) (heart tracing) is also a baseline NCCP test.

STEP 2

Taking or receiving treatment

Swallow tablets whole, with or without food, at about the same time daily. If no more than 12 hours late, take the missed dose when remembered; if more than 12 hours late, skip it and resume the next day. Never double the dose.

STEP 3

Tests and reviews

Repeat blood count, kidney and liver tests and blood glucose as arranged. Bone blood tests and ECG are repeated when clinically indicated. PSA, testosterone and scans follow the individual cancer plan.

STEP 4

Specific instructions

If treatment is interrupted for more than 7 days, contact the team before restarting: a new loading dose is normally needed. Some interacting medicines require a different dose or separation by at least 6 hours. Do not change the dose yourself; ask pharmacy to check every new medicine.

STEP 5

Leaving with a clear plan

Confirm the exact next dose or appointment, duration and verified daytime and out-of-hours contacts.

Sources: NCCP 00830 v3: official regimen. Source reconciled draft; independent clinical, pharmacy and patient/PPI review pending.