Goserelin 10.8 mg: hormone implant every 12 weeks

National Cancer Control Programme (NCCP) 00477 · Source reconciled draft; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Before starting

Full blood count, kidney and liver tests, blood glucose and a bone blood profile. Review heart disease, blood pressure, diabetes, mood, falls and fracture risk, sexual health, medicines and the treatment goal.

STEP 2

Taking or receiving treatment

A trained healthcare professional places the implant under the skin of the front of the abdomen. This 10.8 mg implant has a different schedule from the 3.6 mg formulation; do not swap strengths or appointment intervals.

STEP 3

Tests and reviews

Regular blood count, kidney and liver tests, blood glucose and bone blood profile. NCCP does not specify one interval for these tests. Review symptoms, PSA and testosterone, with scans according to the individual plan.

STEP 4

Specific instructions

Tell the team if you take blood-thinning medicines or are very thin. These can increase bleeding risk at implantation. Severe abdominal pain, marked swelling, dizziness or collapse afterwards needs urgent help. Do not stop blood thinners yourself.

STEP 5

Leaving with a clear plan

Confirm the exact next dose or appointment, duration and verified daytime and out-of-hours contacts.

Sources: NCCP 00477 v3: official regimen. Source reconciled draft; independent clinical, pharmacy and patient/PPI review pending.