Triptorelin 11.25 mg: hormone injection every 12 weeks

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

One injection every 12 weeks

Active medicines: Triptorelin 11.25 mg

Emetogenic risk: Minimal

No routine prophylactic antiemetic order. Agree symptom treatment if needed.

Additional phase-specific support

- A trained healthcare professional prepares and gives the injection using the product-specific instructions. Confirm the 11.25 mg formulation, agreed route and next appointment. Other triptorelin strengths have different schedules.
- Repeat blood count, kidney and liver tests, glucose and bone blood profile when clinically indicated. Review symptoms, PSA and testosterone; scans and the frequency of tests follow the individual plan.
- The team assesses temporary anti-androgen protection against the initial hormone flare. Report worsening pain or trouble passing urine promptly. New weakness or changes in bladder/bowel control require emergency assessment. Keep the written injection dates even if you feel well.
- Bone health, calcium and vitamin D according to individual needs and local policy.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

Sources checked 2026-11-14 · SACTCheck v0.84.52 · Regimen-level drafts; prescriber and oncology-pharmacy verification required