Degarelix: hormone injections every 28 days

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Two starting injections; then every 28 days

Active medicines: Degarelix

Emetogenic risk: Minimal

No routine prophylactic antiemetic order. Agree symptom treatment if needed.

Additional phase-specific support

- A trained healthcare professional gives the injections under the skin of the abdomen. Injection sites are changed and kept away from pressure from belts or waistbands. This formulation is not given into a vein or muscle.
- Repeat blood count, kidney and liver tests, blood glucose and bone blood profile when clinically indicated. Review symptoms and the injection sites. PSA, testosterone and scans follow the individual cancer plan.
- Book the next injection before leaving. If an appointment is missed, contact the team promptly for a new date; do not arrange an extra loading dose yourself. Degarelix does not produce an initial testosterone surge, so anti-androgen treatment solely for flare prevention is not normally needed.
- Bone health, calcium and vitamin D according to individual needs and local policy.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

Sources checked 2026-11-14 · SACTCheck v0.84.52 · Regimen-level drafts; prescriber and oncology-pharmacy verification required