Relugolix: daily hormone treatment

> **Supportive prescription draft for review.** This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Loading dose, then once daily

Active medicines: Relugolix

Emetogenic risk: Minimal to low (oral)

No routine prophylactic antiemetic order. Agree symptom treatment if needed.

Additional phase-specific support

- Swallow tablets whole, with or without food, at about the same time daily. If no more than 12 hours late, take the missed dose when remembered; if more than 12 hours late, skip it and resume the next day. Never double the dose.
- Repeat blood count, kidney and liver tests and blood glucose as arranged. Bone blood tests and electrocardiogram (ECG) are repeated when clinically indicated. PSA, testosterone and scans follow the individual cancer plan.
- If treatment is interrupted for more than 7 days, contact the team before restarting: a new loading dose is normally needed. Some interacting medicines require a different dose or separation by at least 6 hours. Do not change the dose yourself; ask pharmacy to check every new medicine.
- Bone health, calcium and vitamin D according to individual needs and local policy.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

Sources checked 2026-11-14 · SACTCheck v0.84.52 · Regimen-level drafts; prescriber and oncology-pharmacy verification required