Leuprorelin 22.5 mg

Hormone treatment to lower testosterone and help control prostate cancer; your team explains the goal for your setting.

Advanced disease, or selected high-risk disease with radiotherapy · NCCP 00479 v3 · Document revised 14 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Advanced disease, or selected high-risk disease with radiotherapy

My overall treatment

One injection every 12 weeks

84 day view

Treatment timing in this window

Injection / implant visit — Day 1

Medicine continues working — Day 2–84

Treatment

Leuprorelin 22.5 mg is injected under the skin every 12 weeks (84 days) in this NCCP regimen. The prescribed duration depends on whether it supports radiotherapy or treats advanced disease.

Review

PSA means prostate-specific antigen. The team reviews symptoms, PSA, testosterone when needed and the next treatment date.

Your injection interval

Start / visit	Between reviews	Next date
Leuprorelin 22.5 mg is injected under the skin every 12 weeks (84 days) in this NCCP regimen. The prescribed duration depends on whether it supports radiotherapy or treats advanced disease.	The depot keeps working between injections.	Use the exact date agreed with your team.

Your team confirms the active phase and dates. Symptoms may occur during or after treatment; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Bloods and health review** NCCP / product information	Full blood count, kidney and liver tests, blood glucose and a bone blood profile. Review heart disease, blood pressure, diabetes, mood, falls and fracture risk, sexual health, medicines and the treatment goal.
During	**Safety and response** NCCP / individual plan	Repeat blood count, kidney and liver tests, glucose and bone blood profile when clinically indicated. PSA and testosterone assess response. A suspected preparation or administration problem needs review and a testosterone check.

Between treatments

- Hormone effects can occur throughout treatment; there is no guaranteed symptom-free week.
- Confirm the exact next injection date. With radiotherapy, ask when hormone treatment starts and ends and how the radiotherapy dates fit. A low PSA does not mean you should stop injections. Tell the team about worsening pain, trouble passing urine or severe headache with vision changes.

Bring your medicine list and exact appointment dates.

When to get help

Call 112 / **999** for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.


What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Mood, sleep and energy
Leuprorelin 22.5 mg
Report severe low mood, marked fatigue or sleep problems. Ask for support early.

2 Hot flushes and sweating
Leuprorelin 22.5 mg
Hormone suppression can cause hot flushes. Ask about practical measures and treatments if troublesome.

3 Heart and blood pressure
Leuprorelin 22.5 mg
Keep planned checks. Chest pain, fainting or severe breathlessness needs urgent assessment.



4 Weight and blood sugar
Leuprorelin 22.5 mg
Blood glucose, weight and cardiovascular risk need review; suitable activity and diet can help.

5 Injection or implant site
Leuprorelin 22.5 mg
Report worsening redness, swelling, severe pain or a persistent reaction.

6 Sexual health and fertility
Leuprorelin 22.5 mg
Reduced desire and erection difficulties are common. Discuss fertility and contraception; ADT is not reliable contraception.

7 Bones and muscles
Leuprorelin 22.5 mg
Ask about bone assessment, resistance exercise, falls prevention, calcium intake and vitamin D needs.

8 Urine and early symptoms
Leuprorelin 22.5 mg
Report difficulty passing urine or worsening pain. New leg weakness or loss of bladder/ bowel control is an emergency.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call your team promptly for new or worsening symptoms, severe low mood or problems taking treatment. New leg weakness, loss of bladder or bowel control, severe chest pain, severe breathing difficulty, collapse or a seizure needs emergency assessment: call 112/999.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Advanced disease, or selected high-risk disease with radiotherapy · NCCP 00479 v3 · Document revised 14 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Leuprorelin 22.5 mg

One injection every 12 weeks

Phase: _____ Duration: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My injection record

Record actual dates; these spaces do not define treatment duration.

Injection	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team records each injection and the next planned date. The record does not define the total treatment duration. Record any delayed dose or changed plan. An entry does not authorise the next injection. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.