Relugolix

Hormone treatment to lower testosterone and help control prostate cancer; your team explains the goal for your setting.

Advanced hormone-sensitive prostate cancer · NCCP 00830 v3 · Document revised 14 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Advanced hormone-sensitive prostate cancer

My overall treatment

Loading dose, then once daily

28 day reference

Reference window · ongoing treatment

Daily tablets — Day 1–28

Treatment

Relugolix 360 mg on the first day, then 120 mg once daily without planned breaks. Your specialist sets the duration and any prescribed changes.

Review

PSA means prostate-specific antigen. The team reviews symptoms, PSA, testosterone when needed and the next treatment date.

Your daily treatment

Start / visit	Between reviews	Next date
Relugolix 360 mg on the first day, then 120 mg once daily without planned breaks. Your specialist sets the duration and any prescribed changes.	Daily tablets continue.	Use the exact date agreed with your team.

Your team confirms the active phase and dates. Symptoms may occur during or after treatment; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Bloods and health review** NCCP / product information	Full blood count, kidney and liver tests, blood glucose and a bone blood profile. Review heart disease, blood pressure, diabetes, mood, falls and fracture risk, sexual health, medicines and the treatment goal. An electrocardiogram (ECG) (heart tracing) is also a baseline NCCP test.
During	**Safety and response** NCCP / individual plan	Repeat blood count, kidney and liver tests and blood glucose as arranged. Bone blood tests and ECG are repeated when clinically indicated. PSA, testosterone and scans follow the individual cancer plan.

Between treatments

- Hormone effects can occur throughout treatment; there is no guaranteed symptom-free week.
- If treatment is interrupted for more than 7 days, contact the team before restarting: a new loading dose is normally needed. Some interacting medicines require a different dose or separation by at least 6 hours. Do not change the dose yourself; ask pharmacy to check every new medicine.

Bring your medicine list and exact appointment dates.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for


You do not need to know which medicine caused a symptom before calling.

1 Mood, sleep and energy
Relugolix
Report severe low mood, marked fatigue or sleep problems. Ask for support early.

3 Heart and blood pressure
Relugolix
Keep planned checks. Chest pain, fainting or severe breathlessness needs urgent assessment.

5 Tablets and digestion
Relugolix
Report persistent diarrhoea, vomiting or difficulty taking tablets.

7 Bones and muscles
Relugolix
Ask about bone assessment, resistance exercise, falls prevention, calcium intake and vitamin D needs.



2 Hot flushes and sweating
Relugolix
Hormone suppression can cause hot flushes. Ask about practical measures and treatments if troublesome.

4 Weight and blood sugar
Relugolix
Blood glucose, weight and cardiovascular risk need review; suitable activity and diet can help.

6 Sexual health and fertility
Relugolix
Reduced desire and erection difficulties are common. Discuss fertility and contraception; ADT is not reliable contraception.

8 Urine and early symptoms
Relugolix
Report difficulty passing urine or worsening pain. New leg weakness or loss of bladder/ bowel control is an emergency.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Call your team promptly for new or worsening symptoms, severe low mood or problems taking treatment. New leg weakness, loss of bladder or bowel control, severe chest pain, severe breathing difficulty, collapse or a seizure needs emergency assessment: call 112/999.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Advanced hormone-sensitive prostate cancer · NCCP 00830 v3 · Document revised 14 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Relugolix

Loading dose, then once daily

Phase: _____ Continuous treatment: team reviews dose

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My review record

Record actual dates; these spaces do not define treatment duration.

Review	Date / initials

My treatment information

Latest review date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team records each review and the current daily dose. Recording boxes do not imply a fixed course or a rest period. Record any held dose or changed plan. An entry does not authorise a restart; keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.