

**Adjuvant medication prescription for chemotherapy High Risk for nausea and vomiting**

|             |  |  |                             |  |
|-------------|--|--|-----------------------------|--|
| <b>Name</b> |  |  | <b>Allergies</b>            |  |
| <b>DOB</b>  |  |  |                             |  |
| <b>MRN</b>  |  |  | <b>Date of prescription</b> |  |

- Aprepitant 125mg OD PO on day of chemotherapy**  
Directions: Take one capsule one hour prior to chemotherapy. Bring with you to Dunmanway day ward.
- Aprepitant 80mg OD PO on days 2 and 3 of chemotherapy**  
Directions: Take one capsule daily for two days starting the day after chemotherapy.
- Dexamethasone 8mg OD PO on days 2-4 of chemotherapy**  
Directions: Take 8mg daily for three days starting the day after chemotherapy.
- Omeprazole 20mg OD PO on days 1-4 of chemotherapy**  
Directions: Take one capsule daily for four days starting the day of chemotherapy.
- Nystatin 100,000units/mL 1ml QDS PO continuous during chemotherapy**  
Directions: Rinse 1mL around the mouth four times a day after meals during chemotherapy regimen.
- Chlorhexidine mouthwash 10ml QDS PO continuous during chemotherapy**  
Directions: Rinse 10mls around the mouth for at least 1 minute four times a day during chemotherapy regimen. The mouthwash should be expelled from the mouth after rinsing.
- Cyclizine 50mg TDS PO PRN**  
Directions: Take one tablet three times a day as required during chemotherapy regimen.
- Metocloperamide 10mg TDS PO PRN**  
Directions: Take one tablet three times a day as required during chemotherapy regimen.
- Loperamide 4mg STAT then 2mg QDS PO PRN, (max 16mg/day) for chemotherapy-induced diarrhoea**  
Directions: Take two tablets after first episode of diarrhoea, then one after every diarrheal episode

Repeat by: \_\_\_\_ cycles  
(cycle length = \_\_\_\_ weeks)

|                      |  |
|----------------------|--|
| <b>Signed:</b>       |  |
| <b>Name (print):</b> |  |
| <b>IMC number:</b>   |  |

**Adjuvant medication prescription for chemotherapy High Risk for nausea and vomiting**

|             |  |  |                             |  |
|-------------|--|--|-----------------------------|--|
| <b>Name</b> |  |  | <b>Allergies</b>            |  |
| <b>DOB</b>  |  |  |                             |  |
| <b>MRN</b>  |  |  | <b>Date of prescription</b> |  |

- 1. Aprepitant 125mg OD PO on day of chemotherapy**  
Directions: Take one capsule one hour prior to chemotherapy. Bring with you to Dunmanway day ward.
- 2. Aprepitant 80mg OD PO on days 2 and 3 of chemotherapy**  
Directions: Take one capsule daily for two days starting the day after chemotherapy.
- 3. Dexamethasone 8mg OD PO on days 2-4 of chemotherapy**  
Directions: Take 8mg daily for three days starting the day after chemotherapy.
- 4. Omeprazole 20mg OD PO on days 1-4 of chemotherapy**  
Directions: Take one capsule daily for four days starting the day of chemotherapy.
- 5. Nystatin 100,000units/mL 1ml QDS PO continuous during chemotherapy**  
Directions: Rinse 1mL around the mouth four times a day after meals during chemotherapy regimen.
- 6. Chlorhexidine mouthwash 10ml QDS PO continuous during chemotherapy**  
Directions: Rinse 10mls around the mouth for at least 1 minute four times a day during chemotherapy regimen. The mouthwash should be expelled from the mouth after rinsing.
- 7. Cyclizine 50mg TDS PO PRN**  
Directions: Take one tablet three times a day as required during chemotherapy regimen.
- 8. Metocloperamide 10mg TDS PO PRN**  
Directions: Take one tablet three times a day as required during chemotherapy regimen.
- 9. Loperamide 4mg STAT then 2mg QDS PO PRN, (max 16mg/day) for chemotherapy-induced diarrhoea**  
Directions: Take two tablets after first episode of diarrhoea, then one after every diarrheal episode

Repeat by: \_\_\_\_ cycles  
(cycle length = \_\_\_\_ weeks)