SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

Weekly cetuximab + FOLFIRI

Colorectal cancer · NCCP 00328 · version 7 · 14-day cycle

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Patient name	DOB	Hospital no.	Date

Treatment context
This combination is used for RAS wild-type bowel cancer that has spread. Your team explains the aim, alternatives and how benefit will be reviewed.

Schedule
Cetuximab on Day 1 and Day 8. Day 1: irinotecan, folinic acid and fluorouracil, then a 46-hour fluorouracil pump; planned disconnection around Day 3. A repeating 14-day cycle. The number of cycles depends on benefit, side effects and your treatment plan.

1. GENERIC SACT RISKS / DISCUSSION POINTS

- ☐ Purpose, expected benefit and treatment plan explained
- ☐ Treatment may be delayed, modified or stopped because of toxicity
- ☐ Unexpected, rare or serious adverse effects can occur
- ☐ Supportive medicines and urgent contact advice discussed
- ☐ Pregnancy avoidance / contraception / fertility discussed where relevant
- ☐ Bone-marrow suppression / infection risk
- ☐ Anaemia / fatigue and low platelets / bleeding
- ☐ Nausea / vomiting and bowel disturbance
- ☐ Mouth soreness / mucositis and skin changes
- ☐ Allergic / infusion reaction and other organ-specific toxicity

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Cetuximab
EGFR-TARGETED ANTIBODY

- ☐ Rash, dry skin, cracks and sore nails
- ☐ Eye inflammation or corneal injury
- ☐ Low magnesium, calcium or potassium
- ☐ Diarrhoea, dehydration and kidney problems
- ☐ Infusion reaction; rare serious lung or skin effects

Irinotecan
TOPOISOMERASE I CHEMOTHERAPY

- ☐ Early cramps, sweating and diarrhoea
- ☐ Delayed diarrhoea and dehydration
- ☐ Low blood counts / serious infection
- ☐ Nausea, mouth soreness and fatigue
- ☐ Hair thinning or loss
- ☐ Rare serious lung or allergic reactions

Folinic acid
CHEMOTHERAPY MODULATOR

- ☐ Enhances fluorouracil activity and toxicity
- ☐ Rare allergic reaction
- ☐ Diarrhoea / mouth soreness in combination

Fluorouracil
CHEMOTHERAPY

- ☐ Diarrhoea, dehydration and mouth ulcers
- ☐ Low blood counts / infection
- ☐ Hand-foot skin soreness
- ☐ Cardiotoxicity including chest pain / ischaemia
- ☐ Severe early toxicity including DPD-related toxicity
- ☐ Pump / central-line complications

Discuss the treatment aim, alternatives, expected benefit, serious and lasting risks, fertility/pregnancy precautions and the contact plan. Confirm RAS selection for colorectal treatment, infusion premedication, skin care and post-treatment mineral checks. Explain DPD testing and the exact pump schedule. Explain early and delayed diarrhoea separately.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00328 v7.

SACTCheck v0.83.6
Developed by Dr Paul O'Brien