# Weekly cetuximab + irinotecan

NCCP 00330 · Draft — independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

## Infusion visits and care between treatments

Cetuximab on Day 1 and Day 8. Irinotecan is given on Day 1; no home fluorouracil pump. A repeating 14-day cycle. The number of cycles depends on benefit, side effects and your treatment plan.

STEP 1

### Before treatment

Confirm tests, medicines and the exact schedule.

STEP 2

### Prepare for the visit

Arrange transport if premedicines or symptoms affect driving.

STEP 3

### At the day ward

Blood results, symptoms and venous access are checked.

STEP 4

### Infusion and observation

Weekly cetuximab: first infusion about two hours, later doses often one hour. Fortnightly cetuximab: about two hours. Premedicines, review and observation add time. Irinotecan usually runs over 90 minutes.

STEP 5

### At home

Follow the skin-care and/or bowel plan. There is no pump to disconnect.

STEP 6

### Next review

Attend Day 8 for weekly cetuximab between chemotherapy cycles. Confirm the next appointment and follow-up tests.

### ▼ Do I need a port?

A peripheral cannula or existing central line may be used. A port is not automatically needed; the team considers your veins and overall treatment plan.

---

### ▼ If the schedule changes

Ask the unit to confirm the next visit and which medicine is being changed. Do not copy dates from another regimen.

---

Sources: NCCP 00330 v6: official regimen. Draft — independent clinical, pharmacy and patient/PPI review pending.