SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

FOLFOX-4

Colorectal cancer · NCCP 00210 · version 9b · 14-day cycle

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Patient name	DOB	Hospital no.	Date

Treatment context
FOLFOX-4 may be given after surgery for stage II or III colon cancer, or for colorectal cancer that has spread. Your team explains the aim, alternatives and how benefit will be reviewed.

Schedule
Day 1: oxaliplatin and folinic acid, then fluorouracil bolus and a 22-hour infusion. Day 2: further fluorouracil bolus and a second 22-hour infusion; completion around Day 3. A repeating 14-day cycle. For adjuvant FOLFOX-4, NCCP describes 12 cycles; metastatic treatment continues while suitable.

1. GENERIC SACT RISKS / DISCUSSION POINTS

- ☐ Purpose, expected benefit and treatment plan explained
- ☐ Treatment may be delayed, modified or stopped because of toxicity
- ☐ Unexpected, rare or serious adverse effects can occur
- ☐ Supportive medicines and urgent contact advice discussed
- ☐ Pregnancy avoidance / contraception / fertility discussed where relevant
- ☐ Bone-marrow suppression / infection risk
- ☐ Anaemia / fatigue and low platelets / bleeding
- ☐ Nausea / vomiting and bowel disturbance
- ☐ Mouth soreness / mucositis and skin changes
- ☐ Allergic / infusion reaction and other organ-specific toxicity

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Oxaliplatin
PLATINUM CHEMOTHERAPY

- ☐ Cold-triggered tingling / throat discomfort
- ☐ Persistent sensory neuropathy, sometimes lasting after treatment
- ☐ Allergic / infusion reaction
- ☐ Low blood counts and infection
- ☐ Nausea, diarrhoea and fatigue
- ☐ Rare serious lung or blood complications

Folinic acid
CHEMOTHERAPY MODULATOR

- ☐ Enhances fluorouracil activity and toxicity
- ☐ Rare allergic reaction
- ☐ Diarrhoea / mouth soreness in combination

Fluorouracil
CHEMOTHERAPY

- ☐ Diarrhoea, dehydration and mouth ulcers
- ☐ Low blood counts / infection
- ☐ Hand-foot skin soreness
- ☐ Cardiotoxicity including chest pain / ischaemia
- ☐ Severe early toxicity including DPD-related toxicity
- ☐ Pump / central-line complications

Discuss the treatment aim, alternatives, expected benefit, serious and lasting risks, fertility/pregnancy precautions and the contact plan. Explain DPD testing and the exact pump schedule.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00210 v9b.

SACTCheck v0.83.6
Developed by Dr Paul O'Brien