SACTCHECK TREATMENT ASSESSMENT

Cetuximab (7 day) and FOLFIRI (14 day) Therapy

NCCP 00328 | 7
SACTCheck v0.83.6

TUMOUR GROUP	INDICATION	COURSE
Gastrointestinal	RAS wild-type metastatic colorectal cancer.	Advanced disease | q14d

TREATMENT CONTEXT	ASSESSMENT ID	ASSESSMENT TIME
Not entered	JSON-DEMO-001	17/10/2026, 01:40

ENCODED CRITERIA RESULT

1 encoded criterion requires clinical review

Review the highlighted protocol criteria and the current NCCP source before the final treatment decision.

ENTERED VALUES AND ENCODED CRITERIA

Standard assessment

DOMAIN	ENTERED VALUE	ENCODED CRITERION	RESULT
Current stomatitis / mucositis grade	Grade 1 - Mild	<= 1	**Dose-modification pathway**
Weeks delayed for this non-haematological toxicity	1 weeks	<= 2 weeks	**Dose-modification pathway**
Worst stomatitis grade	Grade 3 - Severe	= 3	**Dose-modification pathway**
Current fluorouracil bolus dose level	Level -1	Linked context required	**Additional clinical review/context**
Current fluorouracil infusion dose level	Level -1	Linked context required	**Additional clinical review/context**

COMPONENT-SPECIFIC PROTOCOL ACTIONS

Protocol-level guidance; verify prescribed dose/current level

Bolus 5-FU: Dose level -2 - 260 mg/m^2
After recovery, reduce the affected components by 1 dose level(s). page 5 | Table 5

Infusional 5-FU: Dose level -2 - 1500 mg/m^2
After recovery, reduce the affected components by 1 dose level(s). page 5 | Table 5

UNASSESSED DOMAINS

ECOG performance status, Known hypersensitivity to a regimen medicine, Pregnant, Breastfeeding, Validated tumour RAS status (colorectal indication), Current cetuximab skin toxicity grade, Number of grade >=3 skin episodes, Skin reaction failed to recover within the source withholding period + 30 more

Blank domains were not assumed normal.

CLINICIAN DECISION

Not documented
Not documented

CLINICAL DECISION SUPPORT - NOT TREATMENT CLEARANCE

SACTCheck applies encoded NCCP protocol criteria to clinician-entered information. This output does not constitute treatment clearance and does not replace complete clinical assessment, the current official NCCP protocol, prescribing information, local policy or professional judgement. The responsible oncology clinician retains responsibility for the final treatment decision.