SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Temozolomide for recurrent or progressive glioma

Adults with grade III or IV malignant glioma recurring or progressing after standard therapy under NCCP 00342. · NCCP 00342 · version 4 · 28-day cycle

Patient name	DOB	Hospital no.	Date

Treatment context
Temozolomide is used to help control a high-grade glioma that has returned or progressed after previous treatment. The team considers previous medicines, current symptoms, scans and your priorities when deciding whether it is suitable.

Schedule
Take temozolomide once daily on Days 1–5, then no temozolomide on Days 6–28. Repeat only when your team authorises the next 28-day cycle. The course is reviewed for benefit and toxicity rather than a fixed six-cycle finish.

1. TREATMENT DISCUSSION

- ☐ Purpose, expected benefit and alternatives explained
- ☐ Indication, medicines, sequence and duration confirmed
- ☐ Blood tests and any individually indicated heart checks discussed
- ☐ Support medicines and phase changes explained
- ☐ Fertility, contraception and breastfeeding discussed where relevant
- ☐ Common, serious and lasting drug-specific risks discussed below
- ☐ Urgent symptoms and local contact details confirmed

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Temozolomide
ORAL CHEMOTHERAPY CAPSULE

- ☐ Low blood counts, infection and bleeding
- ☐ Nausea / vomiting and appetite loss
- ☐ Fatigue, headache and constipation
- ☐ Liver injury, including delayed injury
- ☐ PJP and other opportunistic infections
- ☐ Rash and rare severe skin reactions
- ☐ Fertility effects and harm to a pregnancy
- ☐ Rare serious bone-marrow disorders

Discuss treatment intent, evidence and alternatives; phase-specific schedules; blood and organ monitoring; infection, nausea, fertility and pregnancy precautions; practical support and any unlicensed use. Keep the separate radiotherapy consent and medication plans where applicable.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00342 v4.

SACTCheck v0.84.38
Developed by Dr Paul O'Brien