SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Docetaxel 75 mg/m² · every 3 weeks

Source-defined breast, lung, gastric, gynaecological and castration-sensitive prostate cancer (including high-risk nonmetastatic prostate disease) · NCCP 00203 · version 7 · 21-day cycle

Patient name	DOB	Hospital no.	Date

Treatment context
The aim and duration depend on the primary cancer and exact NCCP indication. Confirm the cancer-specific plan before treatment; this shared medicine schedule does not replace that discussion.

Schedule
Docetaxel 75 mg/m² by IV infusion on Day 1 every 21 days. For the source breast indication, continue until progression or unacceptable toxicity; the team reviews benefit and tolerability. The six-cycle source maximum applies to NSCLC.

1. TREATMENT DISCUSSION

- ☐ Purpose, expected benefit and alternatives explained
- ☐ Indication, medicines, sequence and duration confirmed
- ☐ Blood tests and any individually indicated heart checks discussed
- ☐ Support medicines and phase changes explained
- ☐ Fertility, contraception and breastfeeding discussed where relevant
- ☐ Common, serious and lasting drug-specific risks discussed below
- ☐ Urgent symptoms and local contact details confirmed

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Docetaxel
TAXANE CHEMOTHERAPY

- ☐ Low blood counts, infection and sepsis
- ☐ Hair loss, mouth soreness and fatigue
- ☐ Numbness / tingling; sometimes lasting
- ☐ Fluid retention and nail changes
- ☐ Allergic infusion reaction

Discuss treatment aim, alternatives, uncertainty, fertility, contraception, access, phase changes, serious and lasting risks, supportive medicines and the urgent contact plan.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00203 v7.

SACTCheck v0.84.21
Developed by Dr Paul O'Brien