SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

TCHP · subcutaneous trastuzumab

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

HER2-positive neoadjuvant breast cancer · NCCP 00731 · version 2 · 21-day cycle

Patient name	DOB	Hospital no.	Date

Treatment context
Treat HER2-positive early breast cancer as part of a plan intended to reduce the risk of it returning.

Schedule
Six 21-day cycles before surgery: IV pertuzumab, fixed-dose trastuzumab under the skin, IV docetaxel and IV carboplatin. Surgery and pathology guide the postoperative HER2 plan. This is not the combined Phesgo injection.

1. TREATMENT DISCUSSION

- ☐ Purpose, expected benefit and alternatives explained
- ☐ Duration and treatment setting discussed
- ☐ Supportive medicines and urgent contact advice discussed
- ☐ Common, serious and lasting agent-specific risks discussed below
- ☐ Correct HER2 medicine and formulation confirmed
- ☐ Heart and other required monitoring explained
- ☐ Pregnancy / contraception / breastfeeding discussed where relevant
- ☐ Companion cancer medicines reviewed using their own information

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Docetaxel
TAXANE CHEMOTHERAPY

- ☐ Low blood counts, infection and sepsis
- ☐ Hair loss, mouth soreness and fatigue
- ☐ Numbness / tingling; sometimes lasting
- ☐ Fluid retention and nail changes
- ☐ Allergic infusion reaction

Carboplatin
PLATINUM CHEMOTHERAPY

- ☐ Low blood counts, infection and bleeding
- ☐ Nausea, vomiting and fatigue
- ☐ Allergic reaction, sometimes after previous exposure
- ☐ Kidney or electrolyte changes

Trastuzumab
HER2-TARGETED ANTIBODY

- ☐ Heart pumping-function reduction / heart failure
- ☐ Infusion or allergic reaction
- ☐ Diarrhoea and fatigue
- ☐ Rare serious lung effects

Pertuzumab
HER2-TARGETED ANTIBODY

- ☐ Diarrhoea and dehydration
- ☐ Infusion / allergic reaction
- ☐ Heart pumping-function reduction
- ☐ Rash and fatigue

Discuss the treatment aim, alternatives, expected benefit, common and serious effects, fertility and pregnancy precautions, route, monitoring, practical support and the agreed contact plan. The treating team completes the approved individual consent process.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00731 v2.

SACTCheck v0.84.16
Developed by Dr Paul O'Brien