SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Fortnightly cetuximab + modified FOLFOX-6

Advanced / metastatic colorectal cancer · NCCP 00733 · version 2a · 14-day cycle

Patient name	DOB	Hospital no.	Date

Treatment context
The aim is to control advanced bowel cancer and symptoms. The team reviews how well treatment is working and whether side effects remain acceptable; there is no single fixed number of cycles for everyone.

Schedule
Cetuximab and modified FOLFOX6 start on Day 1, followed by one 46-hour fluorouracil infusion with planned disconnection around Day 3. Repeat every 14 days while beneficial and tolerable. No weekly Day 8 cetuximab is scheduled.

1. TREATMENT DISCUSSION

- ☐ Purpose, expected benefit and alternatives explained
- ☐ Indication, medicines, sequence and duration confirmed
- ☐ Blood tests and any individually indicated heart checks discussed
- ☐ DPD result and severe fluoropyrimidine risks reviewed
- ☐ Support medicines and phase changes explained
- ☐ Fertility, contraception and breastfeeding discussed where relevant
- ☐ Common, serious and lasting drug-specific risks discussed below
- ☐ Urgent symptoms and local contact details confirmed

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Cetuximab
EGFR-TARGETED ANTIBODY

- ☐ Rash, dry skin, cracks and sore nails
- ☐ Eye inflammation or corneal injury
- ☐ Low magnesium, calcium or potassium
- ☐ Diarrhoea, dehydration and kidney problems
- ☐ Infusion reaction; rare serious lung or skin effects

Oxaliplatin
PLATINUM CHEMOTHERAPY

- ☐ Cold-triggered tingling / throat discomfort
- ☐ Persistent sensory neuropathy, sometimes lasting after treatment
- ☐ Allergic / infusion reaction
- ☐ Low blood counts and infection
- ☐ Nausea, diarrhoea and fatigue
- ☐ Rare serious lung or blood complications

Folinic acid
CHEMOTHERAPY MODULATOR

- ☐ Enhances fluorouracil activity and toxicity
- ☐ Rare allergic reaction
- ☐ Diarrhoea / mouth soreness in combination

Fluorouracil
CHEMOTHERAPY

- ☐ Diarrhoea, dehydration and mouth ulcers
- ☐ Low blood counts / infection
- ☐ Hand-foot skin soreness
- ☐ Cardiotoxicity including chest pain / ischaemia
- ☐ Severe early toxicity including DPD-related toxicity
- ☐ Pump / central-line complications

Discuss the treatment aim, alternatives, expected benefit, serious and lasting risks, fertility/pregnancy precautions and the contact plan. Confirm RAS selection for colorectal treatment, infusion premedication, skin care and post-treatment mineral checks. Explain DPD testing and the exact pump schedule.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00733 v2a.

SACTCheck v0.84.21
Developed by Dr Paul O'Brien