Pembrolizumab + paclitaxel + carboplatin AUC6

NCCP 00579 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Appointments and time in the unit

Pembrolizumab first over 30 minutes, then paclitaxel over 3 hours, then carboplatin over 30 minutes during induction. Premedication and observation add time. Clinic time includes assessment, preparation and waiting; ask your unit about the full visit.

STEP 2

Treatment phase

Induction: all three medicines on Day 1 every 21 days for up to four cycles. Maintenance: paclitaxel and carboplatin stop; pembrolizumab alone may continue every 21 days while appropriate. The team agrees duration and review points.

STEP 3

Medicines and practical help

Check the paclitaxel premedication and anti-sickness plan for induction. Maintenance has different medicines and may be a shorter visit. Plan transport if medicines make you drowsy. Keep the immunotherapy alert card and urgent numbers with you.

STEP 4

After treatment

Response imaging follows the consultant plan; no fixed universal scan interval is specified in these NCCP regimens. Immune toxicity can arise after the final pembrolizumab dose. Continue symptom vigilance, immunotherapy alert card and the team’s follow-up plan. Review persistent neuropathy, hearing effects and cytopenias even after chemotherapy stops; immunotherapy maintenance has a different supportive-care plan.

Sources: NCCP 00579 v5a · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.