Nivolumab, ipilimumab, pemetrexed and cisplatin

NCCP 00714 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Alternating visits and duration

The first cycle lasts 42 days: all four medicines on Day 1, then nivolumab, pemetrexed and cisplatin on Day 22. Chemotherapy ends after those two visits. From Cycle 2, nivolumab continues on Days 1 and 22 and ipilimumab on Day 1 of each 42-day cycle. Immune treatment continues while appropriate until the cancer progresses, side effects require stopping, or up to 24 months without progression. Plans can change after review.

STEP 2

Time in the Day Ward

Nivolumab over 30 minutes first; ipilimumab over 30 minutes on Day 1 only; pemetrexed over 10 minutes; cisplatin over 60 minutes, starting 30 minutes after pemetrexed, with pre/post hydration. Ipilimumab requires vital signs every 30 minutes during infusion and for one hour afterwards. Clinic time also includes assessment, preparation, observation and support.

STEP 3

Medicine and monitoring appointments

Folic acid starts before pemetrexed, continues through both chemotherapy visits and for 21 days after the last pemetrexed dose. B12 is given in the preceding week. The source repeats B12 every three cycles if pemetrexed continues; this planned course has only two pemetrexed administrations, so do not automatically continue B12 in immune-only cycles. Your team reconciles the steroid used to prevent pemetrexed rash with the anti-sickness steroid into one clear prescription. Do not add the two schedules together or use leftover steroids yourself.

STEP 4

Transport and access

IV treatment through agreed venous access; no home chemotherapy pump. Arrange transport, lab appointments and prescriptions. Allow extra time for pre/post hydration and review fluid restrictions, urine and electrolyte needs individually. Keep immune alert information available for every healthcare contact.

Sources: NCCP 00714 v4 · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.