SACTCHECK · TREATMENT PASSPORT

Nivolumab, ipilimumab, pemetrexed and cisplatin

A record of what you experience

One cycle per page. Each medicine has its own symptom table. Mark the worst symptoms since your last review, when they occurred and what they stopped you doing. Overlapping symptoms can be recorded under more than one medicine; you do not have to identify the cause.

These are patient-language severity prompts, not a validated questionnaire or a formal CTCAE grading tool. Your team assigns the clinical grade. Serious symptoms override the columns: call without waiting to decide a severity level.

The first cycle lasts 42 days: all four medicines on Day 1, then nivolumab, pemetrexed and cisplatin on Day 22. Chemotherapy ends after those two visits. From Cycle 2, nivolumab continues on Days 1 and 22 and ipilimumab on Day 1 of each 42-day cycle. Immune treatment continues while appropriate until the cancer progresses, side effects require stopping, or up to 24 months without progression. Plans can change after review.

The four pages are reusable recording sheets, not a prescribed four-cycle course. Ask for additional sheets as needed.

Emergency: 112 / 999. Severe breathing difficulty, collapse, seizure or severe chest pain needs emergency help.

Contact oncology immediately for temperature ≥37.5°C or ≤36°C, shivering or suddenly feeling unwell, even with a normal temperature. Follow your own hospital’s instructions. If you cannot reach the team, seek urgent medical assessment.

YOUR ONCOLOGY TEAM · DAYTIME

Insert numbers here

Acute Oncology Service

YOUR ONCOLOGY TEAM · OUT OF HOURS

Insert numbers here

Ask for medical oncology

EVENINGS / OVERNIGHT

Your out-of-hours pathway / nearest ED

Other hospitals: use the numbers your team provides.

Your own hospital / contact instructions:

https://sactcheck.com/patient/00714/

Full patient information, sources and visual timeline. No identifiers are encoded.

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

Nivolumab, ipilimumab, pemetrexed and cisplatin

Cycle number: _____ Start date: _____ Next review: _____

Each card represents a 42-day cycle with treatment on Days 1 and 22. Ipilimumab is Day 1 only. Pemetrexed and cisplatin occur only in Cycle 1.

Cycle 1 · Day 1 all four / Day 22 no ipilimumab · copy this sheet for another cycle of the same phase.

Nivolumab · Immunotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Breathing / cough	☐ Usual breathing	☐ New cough or breathlessness: call promptly	☐ Symptoms with usual activity: urgent review	☐ Breathless at rest: urgent assessment	☐ Severe breathing difficulty / collapse: 112/999
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Ipilimumab · Immunotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Skin / nail changes	☐ None	☐ Mild change: report	☐ Pain or daily tasks affected: call	☐ Spreading infection or blistering: urgent review	☐ Severe illness / collapse: emergency help
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Pemetrexed · Chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Mouth soreness / ulcers	☐ None	☐ Mild soreness; eating normally	☐ Painful; soft food needed: call	☐ Cannot eat / drink enough: urgent review	☐ Major breathing difficulty: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999
Skin / nail changes	☐ None	☐ Mild change: report	☐ Pain or daily tasks affected: call	☐ Spreading infection or blistering: urgent review	☐ Severe illness / collapse: emergency help

Cisplatin · Chemotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Numbness / tingling	☐ None	☐ Mild change; usual tasks unaffected	☐ Limits buttons, writing or walking: call	☐ Washing, dressing or walking difficult: urgent review	☐ Sudden major weakness / paralysis: emergency help
Bleeding / bruising	☐ None	☐ New small bruise / brief bleed: report	☐ Repeated bleeding: call today	☐ Bleeding that will not stop: urgent help	☐ Heavy bleeding / collapse: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

> Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

NCCP 00714 V4 · ONE CYCLE, ONE PAGE

Nivolumab, ipilimumab, pemetrexed and cisplatin

Cycle number: _____ Start date: _____ Next review: _____

Each card represents a 42-day cycle with treatment on Days 1 and 22. Ipilimumab is Day 1 only. Pemetrexed and cisplatin occur only in Cycle 1.

Cycle 2 · immune only · ipilimumab Day 1 · copy this sheet for another cycle of the same phase.

Nivolumab · Immunotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Breathing / cough	☐ Usual breathing	☐ New cough or breathlessness: call promptly	☐ Symptoms with usual activity: urgent review	☐ Breathless at rest: urgent assessment	☐ Severe breathing difficulty / collapse: 112/999
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Ipilimumab · Immunotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Skin / nail changes	☐ None	☐ Mild change: report	☐ Pain or daily tasks affected: call	☐ Spreading infection or blistering: urgent review	☐ Severe illness / collapse: emergency help
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

NCCP 00714 V4 · ONE CYCLE, ONE PAGE

Nivolumab, ipilimumab, pemetrexed and cisplatin

Cycle number: _____ Start date: _____ Next review: _____

Each card represents a 42-day cycle with treatment on Days 1 and 22. Ipilimumab is Day 1 only. Pemetrexed and cisplatin occur only in Cycle 1.

Subsequent 42-day cycle · ipilimumab Day 1 · copy this sheet for another cycle of the same phase.

Nivolumab · Immunotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Breathing / cough	☐ Usual breathing	☐ New cough or breathlessness: call promptly	☐ Symptoms with usual activity: urgent review	☐ Breathless at rest: urgent assessment	☐ Severe breathing difficulty / collapse: 112/999
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Ipilimumab · Immunotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Skin / nail changes	☐ None	☐ Mild change: report	☐ Pain or daily tasks affected: call	☐ Spreading infection or blistering: urgent review	☐ Severe illness / collapse: emergency help
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.

NCCP 00714 V4 · ONE CYCLE, ONE PAGE

Nivolumab, ipilimumab, pemetrexed and cisplatin

Cycle number: _____ Start date: _____ Next review: _____

Each card represents a 42-day cycle with treatment on Days 1 and 22. Ipilimumab is Day 1 only. Pemetrexed and cisplatin occur only in Cycle 1.

Subsequent 42-day cycle · repeat if prescribed · copy this sheet for another cycle of the same phase.

Nivolumab · Immunotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Breathing / cough	☐ Usual breathing	☐ New cough or breathlessness: call promptly	☐ Symptoms with usual activity: urgent review	☐ Breathless at rest: urgent assessment	☐ Severe breathing difficulty / collapse: 112/999
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Ipilimumab · Immunotherapy

Symptom	None	Mild	Moderate / call	Severe / urgent	Emergency signs
Diarrhoea	☐ Usual pattern	☐ Up to 3 extra stools above normal	☐ 4–6 extra stools/day or at night: call	☐ 7+ extra stools/day or dehydration: urgent review	☐ Collapse / severe dehydration: 112/999
Skin / nail changes	☐ None	☐ Mild change: report	☐ Pain or daily tasks affected: call	☐ Spreading infection or blistering: urgent review	☐ Severe illness / collapse: emergency help
Fatigue	☐ Usual energy	☐ Tiredness improves with rest	☐ Rest does not help; usual tasks affected: call	☐ Self-care difficult: urgent review	☐ Collapse: 112/999

Dates / doses missed or changed / symptoms affecting daily life / questions:

Red flags override the table. Fever, shivers, suddenly unwell, chest symptoms or inability to drink: seek prompt advice. Severe breathing difficulty, collapse or severe chest pain: 112 / 999.