Durvalumab 1500 mg every 28 days

NCCP 00655 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Two different parts of treatment

Platinum chemotherapy and radiation are given together first. Durvalumab follows after that course has finished and the team has confirmed that the cancer has not progressed.

STEP 2

Infusion visits

The usual reference dose is 1500 mg, given into a vein over about 60 minutes every 28 days. Allow extra time for the clinical review, tests and preparation. If body weight is 30 kg or less, the team must prescribe an appropriate weight-based dose and interval instead of the fixed 1500 mg dose.

STEP 3

Bloods, scans and transport

Confirm where blood tests are taken, who reviews them and the planned clinic/scan appointments. Allow extra time at the treatment unit and arrange help with travel if fatigue or symptoms make it difficult.

STEP 4

A pause or early finish

Treatment can be paused to investigate symptoms or treat an immune reaction. A missed dose does not automatically extend the course beyond the planned maximum. Restart decisions belong to the consultant team.

Sources: NCCP 00655 v3a · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.