SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

Capecitabine

Colorectal and breast cancer · NCCP 00216 · version 8 · 21-day cycle

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Patient name	DOB	Hospital no.	Date

Treatment context
Capecitabine is chemotherapy taken as tablets. NCCP 00216 covers several colorectal and breast cancer settings, including treatment after surgery and treatment of advanced disease. Ask which aim and duration apply to you.

Schedule
capecitabine twice daily on Days 1–14, then no capecitabine on Days 15–21. Follow your individual prescription.

1. GENERIC SACT RISKS / DISCUSSION POINTS

- ☐ Purpose, expected benefit and treatment plan explained
- ☐ Treatment may be delayed, modified or stopped because of toxicity
- ☐ Unexpected, rare or serious adverse effects can occur
- ☐ Supportive medicines and urgent contact advice discussed
- ☐ Pregnancy avoidance / contraception / fertility discussed where relevant
- ☐ Bone-marrow suppression / infection risk
- ☐ Anaemia / fatigue and low platelets / bleeding
- ☐ Nausea / vomiting and bowel disturbance
- ☐ Mouth soreness / mucositis and skin changes
- ☐ Allergic / infusion reaction and other organ-specific toxicity

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Capecitabine

ORAL CHEMOTHERAPY

- ☐ Diarrhoea and dehydration
- ☐ Painful hands / feet
- ☐ Mouth soreness and reduced intake
- ☐ Low blood counts / serious infection
- ☐ Chest pain / heart problems
- ☐ Severe early toxicity, including DPD-related toxicity

Confirm DPD testing, tablet directions, missed-dose and stop-and-call instructions, kidney/liver and blood monitoring, interactions, fertility and contraception. Confirm the indication-specific aim, expected benefit and alternatives.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00216 v8.

SACTCheck v0.83.4
Developed by Dr Paul O'Brien