SACTCHECK TREATMENT ASSESSMENT

Capecitabine + radiotherapy

NCCP 00586 | 3
SACTCheck v0.83.4

TUMOUR GROUP	INDICATION	COURSE
Gastrointestinal	**Locally advanced rectal cancer.**	**Advanced disease | Up to 6 cycles | q7d**

TREATMENT CONTEXT	ASSESSMENT ID	ASSESSMENT TIME
Not entered	**JSON-DEMO-001**	**16/10/2026, 23:35**

ENCODED CRITERIA RESULT

1 encoded criterion requires clinical review

Review the highlighted protocol criteria and the current NCCP source before the final treatment decision.

ENTERED VALUES AND ENCODED CRITERIA

Standard assessment

DOMAIN	ENTERED VALUE	ENCODED CRITERION	RESULT
Creatinine clearance category	30-50 mL/min	>= 30	**Dose-modification pathway**
Creatinine clearance category	30-50 mL/min	<= 50	**Dose-modification pathway**
Haemodialysis	No	= Yes	**Criterion met**

COMPONENT-SPECIFIC PROTOCOL ACTIONS

Protocol-level guidance; verify prescribed dose/current level

Capecitabine: Use 75% of documented starting dose - 75% of documented starting dose per administration; confirm prior reductions and pharmacy tablet banding
Use 75% of the original capecitabine dose for CrCl 30-50 mL/min; apply tablet dose banding and consultant review. page 3 | Table 2

UNASSESSED DOMAINS

ECOG performance status, Hypersensitivity to capecitabine or its excipients, Previous severe, unexpected fluoropyrimidine reaction, Brivudine now or within the previous four weeks, Pregnant, Breastfeeding, DPD status, Absolute neutrophil count + 18 more

Blank domains were not assumed normal.

CLINICIAN DECISION

Not documented
Not documented

CLINICAL DECISION SUPPORT - NOT TREATMENT CLEARANCE

SACTCheck applies encoded NCCP protocol criteria to clinician-entered information. This output does not constitute treatment clearance and does not replace complete clinical assessment, the current official NCCP protocol, prescribing information, local policy or professional judgement. The responsible oncology clinician retains responsibility for the final treatment decision.