SACTCHECK TREATMENT ASSESSMENT

Bevacizumab + capecitabine

NCCP 00623 | 3a
SACTCheck v0.83.4

TUMOUR GROUP	INDICATION	COURSE
Gastrointestinal	**Metastatic or unresectable colorectal cancer when irinotecan- or oxaliplatin-combination chemotherapy is unsuitable.**	**Advanced disease | Until progression/toxicity | q21d**

TREATMENT CONTEXT	ASSESSMENT ID	ASSESSMENT TIME
Not entered	**JSON-DEMO-001**	**16/10/2026, 23:36**

ENCODED CRITERIA RESULT

1 encoded criterion requires clinical review

Review the highlighted protocol criteria and the current NCCP source before the final treatment decision.

ENTERED VALUES AND ENCODED CRITERIA

Standard assessment

DOMAIN	ENTERED VALUE	ENCODED CRITERION	RESULT
24-hour urine total protein	4.1 g/24h	> 4 g/24h	**Criterion not met - urgent review**

COMPONENT-SPECIFIC PROTOCOL ACTIONS

Protocol-level guidance; verify prescribed dose/current level

Bevacizumab: Discontinue - Not to be administered
Discontinue bevacizumab for protein >4 g/24h. page 7 | Table 7

UNASSESSED DOMAINS

ECOG performance status, Hypersensitivity to capecitabine or its excipients, Previous severe, unexpected fluoropyrimidine reaction, Brivudine now or within the previous four weeks, Pregnant, Breastfeeding, DPD status, Assessment timing (optional) + 46 more

Blank domains were not assumed normal.

CLINICIAN DECISION

Not documented
Not documented

CLINICAL DECISION SUPPORT - NOT TREATMENT CLEARANCE

SACTCheck applies encoded NCCP protocol criteria to clinician-entered information. This output does not constitute treatment clearance and does not replace complete clinical assessment, the current official NCCP protocol, prescribing information, local policy or professional judgement. The responsible oncology clinician retains responsibility for the final treatment decision.