BASELINE → TREATMENT → FOLLOW-UP

Checks through your treatment

Baseline preparation, checks before each cycle and tests arranged because of symptoms are different. No fixed scan interval is imposed here.

Check	Before starting	Days 1–5	Days 6–12	Days 13–28	Each next cycle / stopping
Blood count, kidney and liver	Baseline FBC, renal/liver profile	Extra tests if clinically needed	Closer blood-count monitoring with kidney impairment	Review recovery and new-cycle tests	Repeat before each cycle; follow abnormalities
Urine protein	Baseline dipstick	Report urinary changes	Additional testing if needed	Review with next-cycle tests	Dipstick before each cycle
Tablets, symptoms and support	Pharmacy calendar and anti-sickness plan	Check exact AM/PM quantities	No catch-up doses; report symptoms	Rest is still a time to report illness	Reconcile dose history, recovery and support
Treatment benefit	Individual aim and alternatives	Observe symptoms and function	Contact team if worsening	Scans according to Consultant plan	Continue while helpful and tolerated

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00382 v4 · official regimen

THE WHOLE TREATMENT JOURNEY

Trifluridine/tipiracil (Lonsurf) monotherapy: treatment and checks

Previously treated metastatic colorectal cancer

Take the prescribed morning and evening tablets on Days 1–5, have no Lonsurf on Days 6–7, then take morning and evening tablets on Days 8–12. Days 13–28 are tablet-free. A new 28-day cycle starts only when the team authorises it.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

Treatment, tests and reviews across the course

Care / check	Before starting	Days 1–5	Days 6–12	Days 13–28	Each next cycle / stopping
Blood count, kidney and liver NCCP 00382	Baseline FBC, renal/liver profile	Extra tests if clinically needed	Closer blood-count monitoring with kidney impairment	Review recovery and new-cycle tests	Repeat before each cycle; follow abnormalities
Urine protein NCCP 00382	Baseline dipstick	Report urinary changes	Additional testing if needed	Review with next-cycle tests	Dipstick before each cycle
Tablets, symptoms and support NCCP 00382	Pharmacy calendar and anti-sickness plan	Check exact AM/PM quantities	No catch-up doses; report symptoms	Rest is still a time to report illness	Reconcile dose history, recovery and support
Treatment benefit NCCP 00382	Individual aim and alternatives	Observe symptoms and function	Contact team if worsening	Scans according to Consultant plan	Continue while helpful and tolerated

What each check is for

1 Two five-day dosing blocks, then recovery

28-day cycle

This oral chemotherapy aims to control colorectal cancer that has spread after earlier treatments, or when those treatments are unsuitable. It combines two ingredients in one medicine. Your team discusses the likely benefit, uncertainty, alternatives and your priorities.

Ask / act: Confirm the current phase, prescription, tests and support medicines.

Sources: NCCP 00382

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-05 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.34