Nivolumab + ipilimumab · colorectal after chemotherapy

To help control MSI-H/dMMR bowel cancer. After previous fluoropyrimidine-based combination chemotherapy.

MSI-H/dMMR metastatic colorectal cancer after fluoropyrimidine-based combination chemotherapy · NCCP 00551 v7a · Document revised 5 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

MSI-H/dMMR metastatic colorectal cancer after fluoropyrimidine-based combination chemotherapy. Use the written phase plan agreed by your team.

My overall treatment

Cycles 1–4: nivolumab 3 mg/kg, then ipilimumab 1 mg/kg, Day 1 every 21 days.

Then nivolumab alone

Start three weeks after the last combination. 240 mg IV/600 mg under skin every 2 weeks, or 1200 mg under skin every 4 weeks. Team selects the plan.

A typical 21-day cycle

Day 1	Day 2–21
Combination visit: cycles 1–4	Between combination visits

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Baseline checks** Required	MSI/MMR tumour result; blood count, kidney/liver, glucose and thyroid.
Each cycle	**Bloods and symptoms** Required	Blood count, kidney/liver, glucose and thyroid before each cycle in both phases.
As needed	**Other tests** Individual / conditional	Cortisol if indicated; scans and markers on your consultant’s plan.
After	**Immune effects** Required follow-up	Monitoring for at least five months after the last dose; longer if needed.

Between treatments

- New diarrhoea, breathlessness, rash, severe fatigue or other symptoms need prompt advice, even between visits.
- Immune effects may begin after treatment stops. Do not change steroids or restart treatment yourself.

Bring your medicine list and alert card. Confirm the next test, treatment and emergency-contact plan.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital’s temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Hormones / nervous system

Nivolumab + ipilimumab

New severe headache, dizziness, confusion, weakness, thirst or extra urine needs urgent advice; hormone effects may need long-term treatment.

3 Heart

Nivolumab + ipilimumab

Chest pain, racing heartbeat or fainting needs urgent assessment.

5 Kidneys


Nivolumab + ipilimumab

Report much less urine, new swelling or a change in kidney tests.

7 Skin / muscles

Nivolumab + ipilimumab

Spreading rash, blisters, severe weakness or muscle pain needs prompt review.



2 Lungs

Nivolumab + ipilimumab

New cough or breathlessness can signal inflammation or another urgent cause.

4 Liver

Nivolumab + ipilimumab

Yellow eyes/skin or dark urine may signal inflammation; tests also check the liver.

6 Bowel

Nivolumab + ipilimumab

New or persistent diarrhoea, blood in stools or severe tummy pain needs urgent advice.

8 Eyes / vision

Nivolumab + ipilimumab

Report eye pain or new visual change promptly; do not wait for the next cycle.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Contact oncology immediately for new breathlessness, persistent diarrhoea, severe tummy pain, yellow skin or eyes, much less urine, spreading or blistering rash, severe headache, marked weakness, confusion, unusual thirst or passing much more urine. Severe breathing difficulty, chest pain, collapse or seizure needs emergency care. Effects can begin after treatment ends.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

MSI-H/dMMR metastatic colorectal cancer after fluoropyrimidine-based combination chemotherapy · NCCP 00551 v7a · Document revised 5 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Nivolumab + ipilimumab · CRC

4 combination cycles every 21 days, then team-selected nivolumab-only schedule.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678
Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

CUT outer border

BACK / TREATMENT RECORD

My cycle record

Record phase and cycle: combination 1–4, then maintenance.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____
Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.