NCCP 00732 · VERSION 2

Your treatment, explained

Advanced / metastatic colorectal cancer · NCCP 00732 v2 · Document revised 5 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Cetuximab · every two weeks

14-day cycle

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

ACT ON SYMPTOMS, NOT THE CALENDAR

When to get help

Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. New breathing difficulty, chest pain, collapse, severe diarrhoea or inability to drink require prompt assessment. Report eye pain, blurred vision, blistering or infected skin promptly.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.

Open your CUH contact / treatment card

Sources: nccp · hse

Choose a topic to read

YOUR COMPLETE REFERENCE

Read what you need, when you need it.

Start with the overview, explore the cycle, or go straight to a question. Urgent advice is always available.

- How each drug works
- Treatment, recovery and review
- Your 14-day timeline
- Baseline and cycle-specific checks
- Whole treatment, scans and checks
- Tablets, visits and practical planning
- Your treatment aim
- Before the first treatment
- Symptoms in context
- Living with treatment
- Visits and treatment sequence
- When treatment changes or ends
- Pauses, recovery and follow-up
- Supportive medicines
- Understanding the evidence

Questions to ask

Sources

UNDERSTAND YOUR TREATMENT

How your medicines work

The aim is to control advanced bowel cancer and symptoms. Treatment continues while it is helping and side effects remain acceptable; your team reviews the plan with you.

EGFR-TARGETED ANTIBODY

Cetuximab

Erbitux



Blocks growth messages at the cell surface

Cetuximab attaches to EGFR on the outside of cells and blocks some growth messages. In this bowel-cancer plan, validated extended RAS testing helps identify whether treatment is suitable. Healthy skin also uses EGFR, so skin and nail effects need early care.

EGFR receives a growth signal → Cetuximab blocks the receptor → Some growth signals are reduced

What to watch for

Report rash or sore nails early. Tell the nurse immediately about infusion symptoms. Eye pain, new breathing problems or severe diarrhoea need prompt advice. Blood tests check minerals during treatment and for two months afterwards.

Sources: smpc

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

Explore your existing anatomy and side-effect map

REGIMEN-SPECIFIC SIDE EFFECTS

Your anatomical side-effect map

A picture of side effects

Match each symptom box to its numbered body location. Tell your team about new symptoms.

1 Eyes

Cetuximab

Eye pain or new visual change needs prompt advice.

3 Breathing / reaction

Cetuximab

New breathing difficulty requires prompt assessment.

5 Bowel


Treatment

Diarrhoea, increased stoma output or dehydration: call early.

7 Blood / infection

Treatment

Fever, shivers or sudden illness: contact oncology immediately.



2 Blood minerals

Cetuximab

Tests check magnesium and other minerals even when you feel well.

4 Infusion symptoms

Cetuximab

Tell the nurse immediately about itching, swelling or flushing.

6 Skin / nails

Cetuximab

Report painful rash, cracks or infected skin early.

8 Infusion access

Treatment delivery

Pain, swelling, leakage or a loose connection needs immediate advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

THE BIGGER TREATMENT PLAN

Treatment, recovery and review

CLINIC → HOME → REVIEW

A 14-day reference cycle

D1 D14

Cetuximab on Day 1 every 14 days while beneficial and tolerable. No Day 8 cetuximab or home chemotherapy pump is scheduled.

A repeating reference cycle sits within a longer treatment plan. The oncology team chooses the number of cycles and confirms changes; no patient dates are required here.

Sources: nccp

BASELINE → TREATMENT → FOLLOW-UP

When checks happen

The columns show order, not equal periods. There is no fixed number of cycles for these plans. Agree who books each check, when results are reviewed and whom to contact if a check is missed.

Check	Before starting	Early treatment	Repeated treatment	Longer-term review	Pause / completion
Blood count, kidney and liver	Baseline blood count, kidney and liver profile	Review results and symptoms before treatment	Regular tests as arranged by the team	Repeat sooner if unwell or results need follow-up	Follow unresolved abnormalities after treatment
Magnesium, calcium and potassium	Check mineral levels before starting	Repeat periodically during treatment	Keep blood-test and replacement appointments	More checks may be needed if levels fall or diarrhoea develops	Monthly checks for two months after the last antibody dose
Skin, nails, eyes and breathing	Discuss existing skin, eye or lung problems	Start the prescribed skin-care plan; report reactions early	Review skin, nails, diarrhoea and new symptoms	Eye pain, worsening breathing or severe rash needs prompt assessment	Continue care until symptoms and mineral levels settle
Cancer response and treatment choices	Record baseline disease and agree review plan	Bring forward review if symptoms worsen	Scans and other tests follow your consultant's treatment plan	Discuss benefit, side effects and whether the goal has changed	Agree continuation, a break, another treatment or supportive care

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00732 v2 · current official regimen checked 5 November 2026 · EMA: Erbitux product information and patient leaflet

THE WHOLE TREATMENT JOURNEY

Cetuximab · every two weeks: treatment and checks

Advanced / metastatic colorectal cancer

Cetuximab on Day 1 every 14 days while beneficial and tolerable. No Day 8 cetuximab or home chemotherapy pump is scheduled.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

On a small screen, swipe the map sideways. The checks below give the same timing in words.

Treatment, tests and reviews across the course

Care / check	Before starting	Early treatment	Repeated treatment	Longer-term review	Pause / completion
Blood count, kidney and liver NCCP 00732	Baseline blood count, kidney and liver profile	Review results and symptoms before treatment	Regular tests as arranged by the team	Repeat sooner if unwell or results need follow-up	Follow unresolved abnormalities after treatment
Magnesium, calcium and potassium NCCP 00732	Check mineral levels before starting	Repeat periodically during treatment	Keep blood-test and replacement appointments	More checks may be needed if levels fall or diarrhoea develops	Monthly checks for two months after the last antibody dose
Skin, nails, eyes and breathing NCCP 00732	Discuss existing skin, eye or lung problems	Start the prescribed skin-care plan; report reactions early	Review skin, nails, diarrhoea and new symptoms	Eye pain, worsening breathing or severe rash needs prompt assessment	Continue care until symptoms and mineral levels settle
Cancer response and treatment choices NCCP 00732	Record baseline disease and agree review plan	Bring forward review if symptoms worsen	Scans and other tests follow your consultant's treatment plan	Discuss benefit, side effects and whether the goal has changed	Agree continuation, a break, another treatment or supportive care

What each check is for

1

A 14-day reference cycle

Clinic → home → review

The aim is to control advanced bowel cancer and symptoms. Treatment continues while it is helping and side effects remain acceptable; your team reviews the plan with you.

Ask / act: Confirm the current phase, prescription, tests and support medicines.

Sources: NCCP 00732

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-05 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.35

Day 1 → Day 14: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

Cetuximab on Day 1 every 14 days while beneficial and tolerable. No Day 8 cetuximab or home chemotherapy pump is scheduled.

Day of cycle	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Cetuximab infusion	IV													
Skin, nails and minerals	Can build over successive cycles													
Energy and bowel symptoms	Variable timing													

Your planned visits: Cetuximab on Day 1. No home chemotherapy pump. A repeating 14-day cycle. The number of cycles depends on benefit, side effects and your treatment plan.

Care continues: Continue the skin-care plan and mineral checks arranged after cetuximab. Ask which medicines to continue and when your next review is planned.

No guaranteed safe days: An infection, reaction or other urgent problem can occur at any time. The graph does not predict a personal low point.

> **Why these bands stay broad:** NCCP defines the treatment days. Symptom bands are broad educational guidance; reactions can occur on any day, and cumulative effects may continue between cycles. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00732 v2 · current official regimen checked 5 November 2026 ↗

> **There are no guaranteed "safe days".** Fever, shivers or suddenly feeling unwell require immediate contact. Serious symptoms override any timeline or diary.

Read the whole cycle as text

Day 1: Treatment and checks

The team reviews blood tests, symptoms and your prescription before treatment. Premedicines and observation are part of cetuximab visits. Know the contact number and the next planned visit.

Day 2–7: Care between visits

Follow the skin-care or bowel-symptom plan provided. There is no pump to remove. Call early for symptoms; do not wait for the next appointment.

Day 8–14: Recovery and the next review

The team uses symptoms, tests and your priorities to decide what happens next. Bring questions and a symptom record if useful; this is optional.

PRACTICAL TREATMENT GUIDE

Visits, access and care between treatments

Cetuximab on Day 1 every 14 days while beneficial and tolerable. No Day 8 cetuximab or home chemotherapy pump is scheduled.



STEP 1

Before the first visit

Confirm the exact regimen, tests, medicines, emergency number and treatment goal. Ask whether a peripheral cannula or an existing line is appropriate; a port is not automatically required.



STEP 2

At the treatment unit

Fortnightly cetuximab infusion time is set from the dose and safe infusion rate; the first dose may take longer. Blood review, preparation, premedicines and observation add time; infusion time is not the total appointment length.



STEP 3

After the infusion

There is no fluorouracil home pump and no disconnection visit in this regimen.



STEP 4

Between doses

Use the written symptom and support plan; do not wait for the next appointment if urgent symptoms appear.



STEP 5

The next decision

Confirm the next visit after any delay or dose change. Treatment continues only while the oncology team judges benefit and tolerability acceptable. Follow-up mineral checks continue after the final EGFR-antibody dose where applicable.

Transport and daily life

Plan transport if antihistamines, symptoms or other medicines affect driving. Ask for practical, work, nutrition or emotional support.

If treatment changes

Ask which component has been held and which visits remain. A treatment delay does not automatically cancel blood tests, line care or symptom follow-up.

Open practical treatment guide · Printable practical guide

Sources: NCCP 00732 v2 · current official regimen checked 5 November 2026. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.

MAKE THE PLAN PRACTICAL

Support around the medicines

1 Skin, nails and minerals

Start the prescribed skin-care plan early, use moisturiser and sun protection, and report painful or infected skin or nails. Keep mineral blood-test appointments during treatment and for two months after cetuximab.

2 Bowel symptoms and fluids

Follow your written rescue plan. Fever, blood, severe pain, reduced urine or inability to drink needs urgent advice.

3 Plan the visit

Allow for blood tests, review, pharmacy preparation, infusion and observation. A port is not automatically required; ask which access is suitable.

4 Comfort and daily life

Use prescribed anti-sickness medicines, pace activities and ask for nutritional, emotional or practical support.

5 The whole treatment plan

Ask how and when the team will assess benefit, which blood tests continue during a pause, and what happens after the final dose. Mineral checks continue monthly for two months after the last antibody treatment.

6 Recovery and the next dose

A delay or reduction is individual. Tell the team about ongoing side effects and previous dose changes; do not assume a calendar date or a stamped card means that treatment can restart.

Sources: nccp · hse

TREATMENT REFERENCE · 01

Your treatment aim

The setting changes what benefit means.

Why this option?

Cetuximab alone is an option for RAS wild-type bowel cancer that has spread after oxaliplatin and irinotecan have failed, when irinotecan cannot be tolerated. Your team explains the aim, alternatives and how benefit will be reviewed.

Sources: nccp

A plan that can change

This regimen is used to control advanced bowel cancer and symptoms. Ask how benefit will be assessed and whether any local treatment is part of your individual plan. Your preferences and priorities matter.

Sources: nccp

TREATMENT REFERENCE · 02

Before the first treatment

Checks reduce avoidable risk.

Blood tests and medicines

Review blood counts, kidney/liver tests, symptoms, allergies and all medicines and supplements. Tell the team about red-meat allergy or tick bites before cetuximab because these can be associated with infusion-reaction risk.

Sources: nccp

Preparing access

Allow for blood tests, review, pharmacy preparation, infusion and observation. A port is not automatically required; ask which access is suitable.

Sources: nccp

TREATMENT REFERENCE · 03

Symptoms in context

Report what changed and how it affects you.

When to call

New breathing difficulty, chest pain, collapse, severe diarrhoea or inability to drink require prompt assessment. Report eye pain, blurred vision, blistering or infected skin promptly.

Sources: nccp · hse

Bowel symptoms

Follow your written rescue plan. Fever, blood, severe pain, reduced urine or inability to drink needs urgent advice.

Sources: nccp

Skin or nerve changes

Start the prescribed skin-care plan early, use moisturiser and sun protection, and report painful or infected skin or nails. Keep mineral blood-test appointments during treatment and for two months after cetuximab.

Sources: nccp

TREATMENT REFERENCE · 04

Living with treatment

Use the support available.

Food, activity and practical help

Use prescribed anti-sickness medicines, pace activities and ask for nutritional, emotional or practical support.

Sources: nccp · hse

Pregnancy and fertility

Discuss fertility preservation and reproductive plans before treatment. Ask for the contraception and breastfeeding instructions for every medicine; the whole regimen may require longer precautions than one medicine alone.

Sources: nccp

TREATMENT REFERENCE · 05

Visits and treatment sequence

Know which steps happen in the clinic and at home.

The exact delivery plan

Cetuximab on Day 1. No home chemotherapy pump. A repeating 14-day cycle. The number of cycles depends on benefit, side effects and your treatment plan.

Sources: nccp

Radiotherapy relationship

This NCCP regimen does not contain a routine radiotherapy course. If radiation is recommended separately, ask whether it is before, during or after these medicines.

Sources: nccp

TREATMENT REFERENCE · 06

When treatment changes or ends

Follow-up continues.

Symptoms and monitoring

Continue the cetuximab skin plan and mineral checks for two months after the last dose. Ask which medicines to continue, when to seek help and how response will be assessed.

Sources: nccp

Next decisions

The team reviews benefit, side effects and your priorities. Supportive and palliative care can help alongside active treatment. Ask for a clear next-step and contact plan.

Sources: nccp · hse

TREATMENT REFERENCE · 07

Pauses, recovery and follow-up

Your team may pause or change one or more medicines. Do not restart or change doses yourself.

Decisions before restarting

The team checks the severity and recovery of side effects, previous episodes and the doses you actually received. A standard table dose does not automatically override a prior individual reduction.

Sources: nccp

Keep testing appointments

Blood tests may continue during a treatment pause. Keep monthly electrolyte, magnesium and calcium checks for two months after the last cetuximab dose. Scans and tumour-marker timing follow the consultant plan; these documents do not impose a universal scan interval.

Sources: nccp

CO.17: KRAS exon 2 analysis

Studies can help explain why a treatment is offered. The schedule, previous medicines and tumour tests affect how well the evidence fits you.

Cetuximab improved cancer control and survival in a historical trial of selected people whose bowel cancer had already been treated. The study used weekly treatment and older tumour testing, so it does not provide an exact personal estimate for this fortnightly plan.

Sources: trial

PREPARE FOR THE CONVERSATION

Questions worth asking

How is my treatment chosen?

Cetuximab alone is an option for RAS wild-type bowel cancer that has spread after oxaliplatin and irinotecan have failed, when irinotecan cannot be tolerated. Your team explains the aim, alternatives and how benefit will be reviewed.

Do I need to enter dates to understand the cycle?

No. The graph explains a general pattern. Your hospital arranges appointments; the guide and its safety information are available without entering anything.

Will I have a pump at home?

There is no home fluorouracil pump in this regimen. Your team chooses a cannula or central line based on your veins and wider treatment plan.

What happens if a dose is delayed?

The team checks the cause and may delay, reduce, change or stop one medicine. Ask which component changes and when you should next attend; do not restart yourself.

How will I know whether it is helping?

Scans, symptoms and clinical review assess benefit. A rash, side effect or good day alone cannot show whether the cancer is responding.

What about radiotherapy?

This NCCP regimen does not contain a routine radiotherapy course. If radiation is recommended separately, ask whether it is before, during or after these medicines.

What about fertility, pregnancy and breastfeeding?

Discuss fertility preservation and reproductive plans before treatment. Ask for the contraception and breastfeeding instructions for every medicine; the whole regimen may require longer precautions than one medicine alone.

Why is RAS testing mentioned?

RAS testing selects cetuximab treatment in colorectal cancer. It is not the selection test for the separate head-and-neck radiotherapy indication. Confirm the cancer-specific plan with your team.

> **Try explaining the plan back:** Can you explain which visits are planned, whether a pump is involved, and which symptoms mean you should call today? No dates or symptom scores are needed to use this guide.

SOURCE TRANSPARENCY

Where this information comes from

Regimen schedule: NCCP 00732 v2. Medicine explanations and safety information: sources below. The timeline is an educational illustration, not a measured risk curve or an individual prediction.

- NCCP 00732 v2 · current official regimen checked 5 November 2026
- HSE: chemotherapy side effects and urgent advice
- EMA: Erbitux product information and patient leaflet
- CO.17 KRAS primary analysis
- NCCP NCIS supportive antiemetic medicines V8 (2025)

SACTCheck · Developed by Dr Paul O'Brien · Sources checked 2026-11-05.
This information supports your treatment discussion. Your oncology team confirms your personal treatment and urgent-care plan.

Written source addresses

https://healthservice.hse.ie/documents/6602/732_v2_Cetuximab_14day.pdf

https://www.ema.europa.eu/en/documents/product-information/erbitux-epar-product-information_en.pdf

https://www2.hse.ie/conditions/chemotherapy/side-effects/

https://pubmed.ncbi.nlm.nih.gov/18946061/

https://healthservice.hse.ie/documents/7180/Supportive_Care_Antiemetics_for_inclusion_NCIS_Medical_Oncology.pdf