NCCP 00207 · VERSION 6

Your treatment, explained

Selected metastatic colorectal cancer; or locally advanced head/neck cancer with radiotherapy when platinum is unsuitable · NCCP 00207 v6 · Document revised 5 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Weekly cetuximab

7-day cycle

Source reconciled; independent clinical, pharmacy and patient/PPI review pending

ACT ON SYMPTOMS, NOT THE CALENDAR

When to get help

Emergency: call 112 / 999

Severe breathing difficulty, collapse, seizure or severe chest pain: call emergency services. Do not wait for an online answer.

Contact your oncology service immediately

Temperature 37.5°C or higher, 36°C or lower, shivering or suddenly feeling unwell — even with a normal temperature. New breathing difficulty, chest pain, collapse, severe diarrhoea or inability to drink require prompt assessment. Report eye pain, blurred vision, blistering or infected skin promptly.

Follow the urgent contact instructions from your treating hospital. If you cannot reach the team, seek urgent medical assessment. Outside clinic hours, use your instructed out-of-hours pathway or the emergency department.

This page does not diagnose symptoms or decide whether it is safe to stay at home. You do not need to know which medicine caused a symptom before calling.

Open your CUH contact / treatment card

Sources: nccp · hse

Choose a topic to read

YOUR COMPLETE REFERENCE

Read what you need, when you need it.

Start with the overview, explore the cycle, or go straight to a question. Urgent advice is always available.

- How each drug works
- Treatment, recovery and review
- Your 7-day timeline
- Baseline and cycle-specific checks
- Whole treatment, scans and checks
- Chemotherapy and radiotherapy timing
- Tablets, visits and practical planning
- Your treatment aim
- Before the first treatment
- Symptoms in context
- Living with treatment
- Visits and treatment sequence
- When treatment changes or ends
- Supportive medicines
- Understanding the evidence
- Questions to ask
- Sources

UNDERSTAND YOUR TREATMENT

How your medicines work

For selected bowel cancer that has spread, cetuximab aims to control the cancer after previous treatments. For selected locally advanced head-and-neck cancer when platinum chemotherapy is unsuitable, it is combined with radiotherapy. Your team confirms which pathway applies.

EGFR-TARGETED ANTIBODY

Cetuximab

Erbitux



Blocks growth messages at the cell surface

Cetuximab attaches to EGFR on the outside of cells and blocks growth messages. In bowel cancer, a validated RAS wild-type result helps identify whether this approach is suitable. In head-and-neck cancer, weekly cetuximab may be combined with radiotherapy under a separate treatment plan. Healthy skin also uses EGFR, so skin and nail effects need early care.

EGFR receives a growth signal → Cetuximab blocks the receptor → Some growth signals are reduced

What to watch for

Report rash or sore nails early. Tell the nurse immediately about infusion symptoms. Eye pain, new breathing problems or severe diarrhoea need prompt advice. Blood tests check minerals during treatment and for two months afterwards.

Sources: smpc

The drawings are simplified explanations of drug action, not anatomical pictures or a prediction of response.

Explore your existing anatomy and side-effect map

REGIMEN-SPECIFIC SIDE EFFECTS

Your anatomical side-effect map

A picture of side effects

Match each symptom box to its numbered body location. Tell your team about new symptoms.

1 Eyes

Cetuximab

Eye pain or new visual change needs prompt advice.

3 Breathing / reaction

Cetuximab

New breathing difficulty requires prompt assessment.

5 Bowel


Treatment

Diarrhoea, increased stoma output or dehydration: call early.

7 Blood / infection

Treatment

Fever, shivers or sudden illness: contact oncology immediately.



2 Blood minerals

Cetuximab

Tests check magnesium and other minerals even when you feel well.

4 Infusion symptoms

Cetuximab

Tell the nurse immediately about itching, swelling or flushing.

6 Skin / nails

Cetuximab

Report painful rash, cracks or infected skin early.

8 Infusion access

Treatment delivery

Pain, swelling, leakage or a loose connection needs immediate advice.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

THE BIGGER TREATMENT PLAN

Treatment, recovery and review

CLINIC → HOME → REVIEW

A 7-day reference cycle

D1 D7

Cetuximab on Day 1. No home chemotherapy pump. A repeating 7-day cycle. The number of cycles depends on benefit, side effects and your treatment plan. For head-and-neck radiotherapy: start cetuximab one week before radiation, then continue weekly during radiation until it ends.


This is a general explanation, not an appointment calendar. Delays or changes are decided by your team. Each medicine can need a different change.

Sources: nccp

THE WHOLE TREATMENT COURSE

Head-and-neck indication: cetuximab before and during radiotherapy

This panel applies only to the head-and-neck radiotherapy indication of NCCP 00207. Cetuximab is a targeted antibody. Colorectal cetuximab alone has no routine radiotherapy component.



The axis shows order, not equal durations. Cetuximab starts one week before radiation and continues until radiation ends. The radiation team sets the number of sessions and confirms eligibility.

> **After the course: continue skin, mouth, swallowing and nutrition support. Mineral monitoring continues for two months after the final cetuximab dose.**

Educational reference, not a personal appointment calendar. NCCP 00207 v5, page 1; current EMA Erbitux information

BASELINE → TREATMENT → FOLLOW-UP

Treatment checks across the course

Relative stages, not equally spaced dates. Agree who books each check, reviews results and gives advice if a check is missed.

Check	Before starting	First treatment	During treatment	Phase / dose review	After treatment
Bloods and symptoms	Baseline blood count, kidney and liver checks	Confirm tests and prescription before the first dose	Repeat bloods and symptom checks according to this regimen	Recheck after a hold or treatment change	Follow unresolved effects and the agreed disease-review plan
Treatment review	Confirm exact cancer setting and biomarkers	Explain expected benefit and how it will be assessed	Report new symptoms between visits	Review response, tolerability and active medicines	Agree the next treatment or surveillance plan
Electrolytes and EGFR effects	Check electrolyte and skin-risk plan	Skin care and prompt reaction reporting	Monitor magnesium, calcium, potassium, skin and eyes	Do not stop follow-up because the infusion stops	Monthly electrolytes, magnesium and calcium for two months after the last EGFR-antibody dose

The team arranges and reviews tests. This guide shows expected timing; it does not record completed appointments. NCCP 00207 v6 · official regimen

THE WHOLE TREATMENT JOURNEY

Weekly cetuximab: treatment and checks

Selected metastatic colorectal cancer; or locally advanced head/neck cancer with radiotherapy when platinum is unsuitable

One larger first infusion is followed by weekly infusions. Bowel-cancer treatment continues while helping and tolerated. For head-and-neck treatment, cetuximab starts one week before radiotherapy and continues until radiation treatment finishes.

Read from left to right. Columns show stages, not equal lengths of time. This is a general plan: no dates or personal information are needed. Your team confirms which steps apply and arranges appointments.

On a small screen, swipe the map sideways. The checks below give the same timing in words.

Treatment, tests and reviews across the course

Care / check	Before starting	First treatment	During treatment	Phase / dose review	After treatment
Bloods and symptoms NCCP 00207	Baseline blood count, kidney and liver checks	Confirm tests and prescription before the first dose	Repeat bloods and symptom checks according to this regimen	Recheck after a hold or treatment change	Follow unresolved effects and the agreed disease-review plan
Treatment review NCCP 00207	Confirm exact cancer setting and biomarkers	Explain expected benefit and how it will be assessed	Report new symptoms between visits	Review response, tolerability and active medicines	Agree the next treatment or surveillance plan

Care / check	Before starting	First treatment	During treatment	Phase / dose review	After treatment
Electrolytes and EGFR effects NCCP 00207	Check electrolyte and skin-risk plan	Skin care and prompt reaction reporting	Monitor magnesium, calcium, potassium, skin and eyes	Do not stop follow-up because the infusion stops	Monthly electrolytes, magnesium and calcium for two months after the last EGFR-antibody dose

What each check is for

1

A 7-day reference cycle

Clinic → home → review

For selected bowel cancer that has spread, cetuximab aims to control the cancer after previous treatments. For selected locally advanced head-and-neck cancer when platinum chemotherapy is unsuitable, it is combined with radiotherapy. Your team confirms which pathway applies.

Ask / act: Confirm the current phase, prescription, tests and support medicines.

Sources: NCCP 00207

At the next review, ask

- Which treatment phase am I in, and what happens next?
- Which test is expected next, who arranges it, and who explains the result?
- Who should I contact if an expected appointment has not arrived?

This map does not show whether a test has been ordered, booked or completed. Do not wait for a scheduled test if you develop new symptoms.

Sources checked 2026-11-05 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending · SACTCheck v0.84.34

Day 1 → Day 7: a general pattern

This is a visual orientation, not a prediction of your personal cycle. It shows when some effects often emerge and which effects are **not** tied to a particular day.

One larger first infusion is followed by weekly infusions. Bowel-cancer treatment continues while helping and tolerated. For head-and-neck treatment, cetuximab starts one week before radiotherapy and continues until radiation treatment finishes.

Day of cycle	1	2	3	4	5	6	7
Cetuximab infusion	IV						
Skin, nails and minerals	Can build over successive cycles						
Energy and bowel symptoms	Variable timing						

Your planned visits: Cetuximab on Day 1. No home chemotherapy pump. A repeating 7-day cycle. The number of cycles depends on benefit, side effects and your treatment plan. For head-and-neck radiotherapy: start cetuximab one week before radiation, then continue weekly during radiation until it ends.

Care continues: Continue the skin-care plan and mineral checks arranged after cetuximab. Ask which medicines to continue and when your next review is planned.

No guaranteed safe days: An infection, reaction or other urgent problem can occur at any time. The graph does not predict a personal low point.

> **Why these bands stay broad:** NCCP defines the treatment days. Symptom bands are broad educational guidance; reactions can occur on any day, and cumulative effects may continue between cycles. Band length does not show likelihood or severity, and the chart cannot tell you whether it is safe to receive treatment. NCCP 00207 v6 · official regimen ↗

> **There are no guaranteed "safe days".** Fever, shivers or suddenly feeling unwell require immediate contact. Serious symptoms override any timeline or diary.

Read the whole cycle as text

Day 1: Treatment and checks

The team reviews blood tests, symptoms and your prescription before treatment. Premedicines and observation are part of cetuximab visits. Know the contact number and the next planned visit.

Day 2–3: Care between visits

Follow the skin-care or bowel-symptom plan provided. There is no pump to remove. Call early for symptoms; do not wait for the next appointment.

Day 4–7: Recovery and the next review

The team uses symptoms, tests and your priorities to decide what happens next. Bring questions and a symptom record if useful; this is optional.

PRACTICAL TREATMENT GUIDE

Infusion visits and care between treatments

Cetuximab on Day 1. No home chemotherapy pump. A repeating 7-day cycle. The number of cycles depends on benefit, side effects and your treatment plan. For head-and-neck radiotherapy: start cetuximab one week before radiation, then continue weekly during radiation until it ends.

STEP 1

Treatment phases and appointments

One larger first infusion is followed by weekly infusions. Bowel-cancer treatment continues while helping and tolerated. For head-and-neck treatment, cetuximab starts one week before radiotherapy and continues until radiation treatment finishes.

STEP 2

Checks after treatment

Monthly electrolytes, magnesium and calcium for two months after the last cetuximab dose; assign responsibility before treatment ends. Response assessment follows the consultant plan; head/neck radiation follow-up is coordinated separately.

STEP 3

Access and practical support

Day-unit IV access; no home fluorouracil pump. Head/neck pathway requires a separate radiation appointment and nutrition/swallowing support plan.

Do I need a port?

A peripheral cannula or existing central line may be used. A port is not automatically needed; the team considers your veins and overall treatment plan.

If the schedule changes

Ask the unit to confirm the next visit and which medicine is being changed. Do not copy dates from another regimen.

Open practical treatment guide · Printable practical guide

Sources: NCCP 00207 v6 · official regimen. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.

MAKE THE PLAN PRACTICAL

Support around the medicines

1 First cetuximab dose

Skin care and sun protection under local EGFR policy. First two doses require post-infusion observation; delayed-reaction education. Monthly electrolytes/Mg/Ca for two months after last dose.

2 Subsequent weekly cetuximab

Premedication continues for subsequent infusions. Head/neck RT requires separate mucosal, skin, swallow and nutrition support. Post-treatment monthly electrolytes/Mg/Ca for two months.

Sources: nccp · hse

TREATMENT REFERENCE · 01

Your treatment aim

The setting changes what benefit means.

Why this option?

Cetuximab alone is an option for RAS wild-type bowel cancer that has spread after oxaliplatin and irinotecan have failed, when irinotecan cannot be tolerated. A separate indication combines weekly cetuximab with radiotherapy for locally advanced head-and-neck cancer; the radiation team confirms suitability, including the current NCCP catalogue criteria. Your team explains the aim, alternatives and how benefit will be reviewed.

Sources: nccp

A plan that can change

For bowel cancer, discuss control of cancer and symptoms, and whether further local treatment may become possible. For head-and-neck treatment, discuss the aim of the combined radiation plan. Your preferences and priorities matter.

Sources: nccp

Separate head-and-neck evidence

A randomised trial compared radiotherapy plus weekly cetuximab with radiotherapy alone and found improved local control and survival. It did not compare cetuximab with cisplatin; it cannot establish that cetuximab is equivalent to cisplatin-based chemoradiotherapy.

Sources: hntrial

TREATMENT REFERENCE · 02

Before the first treatment

Checks reduce avoidable risk.

Blood tests and medicines

Review blood counts, kidney/liver tests, symptoms, allergies and all medicines and supplements. Tell the team about red-meat allergy or tick bites before cetuximab because these can be associated with infusion-reaction risk.

Sources: nccp

Preparing access

Allow for blood tests, review, pharmacy preparation, infusion and observation. A port is not automatically required; ask which access is suitable.

Sources: nccp

TREATMENT REFERENCE · 03

Symptoms in context

Report what changed and how it affects you.

When to call

New breathing difficulty, chest pain, collapse, severe diarrhoea or inability to drink require prompt assessment. Report eye pain, blurred vision, blistering or infected skin promptly.

Sources: nccp · hse

Bowel symptoms

Follow your written rescue plan. Fever, blood, severe pain, reduced urine or inability to drink needs urgent advice.

Sources: nccp

Skin and nail changes

Start the prescribed skin-care plan early, use moisturiser and sun protection, and report painful or infected skin or nails. Keep mineral blood-test appointments during treatment and for two months after cetuximab.

Sources: nccp

TREATMENT REFERENCE · 04

Living with treatment

Use the support available.

Food, activity and practical help

Use prescribed anti-sickness medicines, pace activities and ask for nutritional, emotional or practical support.

Sources: nccp · hse

Pregnancy and fertility

Discuss fertility preservation and reproductive plans before treatment. Ask for the contraception and breastfeeding instructions for every medicine; the whole regimen may require longer precautions than one medicine alone.

Sources: nccp

TREATMENT REFERENCE · 05

Visits and treatment sequence

Know which steps happen in the clinic and at home.

The exact delivery plan

Cetuximab on Day 1. No home chemotherapy pump. A repeating 7-day cycle. The number of cycles depends on benefit, side effects and your treatment plan. For head-and-neck radiotherapy: start cetuximab one week before radiation, then continue weekly during radiation until it ends.

Sources: nccp

Radiotherapy relationship

For the head-and-neck indication, cetuximab starts one week before radiation and continues weekly alongside it. The graphic below shows the relationship. Colorectal cetuximab alone does not include a routine radiation course.

Sources: nccp

Head-and-neck radiation support

Ask the radiation team about dental review, mouth care, pain relief, nutrition and swallowing support. Radiation-related mouth/throat soreness and skin changes can build during the course and continue afterwards. Trouble swallowing fluids or worsening breathing needs urgent assessment.

Sources: smpc

TREATMENT REFERENCE · 06

When treatment changes or ends

Follow-up continues.

Symptoms and monitoring

Continue the cetuximab skin plan and mineral checks for two months after the last dose. Ask which medicines to continue, when to seek help and how response will be assessed.

Sources: nccp

Next decisions

The team reviews benefit, side effects and your priorities. Supportive and palliative care can help alongside active treatment. Ask for a clear next-step and contact plan.

Sources: nccp · hse

Why this treatment is offered

In selected previously treated bowel cancer, cetuximab improved disease control and survival compared with supportive care. Separate head-and-neck evidence supports adding it to radiotherapy in the appropriate setting.

In selected previously treated bowel cancer, cetuximab improved disease control and survival compared with supportive care. Separate head-and-neck evidence supports adding it to radiotherapy in the appropriate setting.

Sources: trial-v08421

PREPARE FOR THE CONVERSATION

Questions worth asking

How is my treatment chosen?

Cetuximab alone is an option for RAS wild-type bowel cancer that has spread after oxaliplatin and irinotecan have failed, when irinotecan cannot be tolerated. A separate indication combines weekly cetuximab with radiotherapy for locally advanced head-and-neck cancer; the radiation team confirms suitability, including the current NCCP catalogue criteria. Your team explains the aim, alternatives and how benefit will be reviewed.

Do I need to enter dates to understand the cycle?

No. The graph explains a general pattern. Your hospital arranges appointments; the guide and its safety information are available without entering anything.

Will I have a pump at home?

There is no home fluorouracil pump in this regimen. Your team chooses a cannula or central line based on your veins and wider treatment plan.

What happens if a dose is delayed?

The team checks the cause and may delay, reduce, change or stop one medicine. Ask which component changes and when you should next attend; do not restart yourself.

How will I know whether it is helping?

Scans, symptoms and clinical review assess benefit. A rash, side effect or good day alone cannot show whether the cancer is responding.

What about radiotherapy?

For the head-and-neck indication, cetuximab starts one week before radiation and continues weekly alongside it. The graphic below shows the relationship. Colorectal cetuximab alone does not include a routine radiation course.

What about fertility, pregnancy and breastfeeding?

Discuss fertility preservation and reproductive plans before treatment. Ask for the contraception and breastfeeding instructions for every medicine; the whole regimen may require longer precautions than one medicine alone.

Why is RAS testing mentioned?

RAS testing selects cetuximab treatment in colorectal cancer. It is not the selection test for the separate head-and-neck radiotherapy indication. Confirm the cancer-specific plan with your team.

Which checks continue when treatment changes or finishes?

Monthly electrolytes, magnesium and calcium for two months after the last cetuximab dose; assign responsibility before treatment ends. Response assessment follows the consultant plan; head/neck radiation follow-up is coordinated separately.

> **Try explaining the plan back:** Can you explain which visits are planned, whether a pump is involved, and which symptoms mean you should call today? No dates or symptom scores are needed to use this guide.

SOURCE TRANSPARENCY

Where this information comes from

Regimen schedule: NCCP 00207 v6. Medicine explanations and safety information: sources below. The timeline is an educational illustration, not a measured risk curve or an individual prediction.

- NCCP 00207 v6 · official regimen
- HSE: chemotherapy side effects and urgent advice
- EMA: Erbitux product information and patient leaflet
- CO.17 KRAS analysis — primary report
- Bonner: radiotherapy with cetuximab, head-and-neck trial
- CO.17 KRAS subgroup; Bonner radiotherapy trial
- NCCP emetogenic classification V6
- NCCP NCIS Medical Oncology antiemetics V8
- Bonner: five-year cetuximab plus radiotherapy follow-up

SACTCheck · Developed by Dr Paul O'Brien · Sources checked 2026-11-05.
This information supports your treatment discussion. Your oncology team confirms your personal treatment and urgent-care plan.

Written source addresses

https://assets.hse.ie/media/documents/207_v5_Cetuximab_7day.pdf

https://healthservice.hse.ie/documents/6941/207_Cetuximab_7day_.pdf

https://www2.hse.ie/conditions/chemotherapy/side-effects/

https://www.ema.europa.eu/en/documents/product-information/erbitux-epar-product-information_en.pdf

https://pubmed.ncbi.nlm.nih.gov/18946061/

https://pubmed.ncbi.nlm.nih.gov/16467544/

https://healthservice.hse.ie/documents/7152/Classification_Document_for_Systemic_Anti-_Cancer_Therapy_SACT_Induced_Nausea__pWgDCrm.pdf

https://healthservice.hse.ie/documents/7180/Supportive_Care_Antiemetics_for_inclusion_NCIS_Medical_Oncology.pdf

https://pubmed.ncbi.nlm.nih.gov/19897418/