Regorafenib (Stivarga) monotherapy

NCCP 00244 · Source reconciled; independent clinical, pharmacy and patient/PPI review pending

PRACTICAL TREATMENT GUIDE

Practical treatment planning

Agree the arrangements with your oncology team.

STEP 1

Prescription and tablet calendar

Confirm the exact daily tablet number, Days 1–21 and Days 22–28, blood-test appointments and who authorises the next cycle. A clinician may prescribe a lower starting dose or change it after side effects; follow the issued prescription.

STEP 2

Taking tablets with food

Swallow the prescribed tablets whole with water at the same time each day after a light meal containing less than 30% fat. Ask the pharmacist for a practical meal example. Follow your own tablet label and calendar; do not change the dose yourself. Avoid grapefruit juice.

STEP 3

Missed or vomited doses

If you miss a dose, take it as soon as you remember on the same day. Do not take two doses in one day or make up the dose the next day. If you vomit after taking regorafenib, do not take extra tablets; tell the team. If unsure, contact the pharmacist or oncology team.

STEP 4

Storage and handling

Keep tablets in their original bottle, tightly closed, away from children and protected from moisture. Record when the bottle is first opened; the leaflet says to discard remaining tablets seven weeks after opening. Ask the pharmacist about safe disposal and help if you cannot swallow whole tablets.

STEP 5

Accidental extra tablets

Seek medical advice straight away if you take more than prescribed. Take the pack and current calendar with you.

STEP 6

Visits and practical support

This is tablet treatment without an infusion or take-home pump, but it still needs regular visits and tests. Ask who arranges appointments and results, and tell the team about travel, work or caring needs before changing the schedule.

Sources: NCCP 00244 v5 · Regorafenib Monotherapy · revised 25 March 2025. Source reconciled; independent clinical, pharmacy and patient/PPI review pending.