Fruquintinib (Fruzaqla) monotherapy

Targeted capsule treatment to help control metastatic bowel cancer after previous therapies.

Metastatic colorectal cancer after standard therapies and trifluridine/tipiracil or regorafenib · NCCP 00890 v2a · Document revised 5 November 2026

Your personal plan: this is the standard regimen schedule. Your team confirms your prescribed doses, cycles, treatment changes and appointments in your written plan.

Previously treated metastatic colorectal cancer; follow the individual prescription and monitoring plan.

My overall treatment

Days 1–21: prescribed dose once daily, with or without food. Days 22–28: no fruquintinib.

28-day cycle
Repeat only when authorised. No fixed final cycle; the team reviews benefit and side effects.

A typical 28-day cycle

Day 1–7	Day 8–21	Day 22–28
Start with a clear calendar	Continue capsules and monitoring	Capsule-free week

Timing may change after review. Side effects can occur on any day; there are no guaranteed safe days.

My tests and checks

When	What is checked	Timing / reason
Before	**Baseline checks** Required	Blood count, kidneys/liver, thyroid, urine protein and controlled blood pressure.
Early / later	**Blood pressure** Required	Weekly for the first month, then monthly or more often if needed.
During	**Bloods and urine** Required / conditional	Regular blood count, kidney/liver, thyroid and urine tests. Warfarin needs extra INR checks.
Review / after	**Response and recovery** Individual plan	Scans, markers and follow-up on your consultant’s plan; review unresolved effects.

Between treatments

- Report painful hands/feet, bleeding, bowel changes or poor intake early, including during the rest week.
- Do not replace a vomited dose. Follow the missed-dose plan; if exactly 12 hours late or unsure, call the team.

Confirm capsule strength, calendar, skin care, test dates and urgent contacts. Swallow capsules whole; do not open them.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital’s temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

What to watch for

You do not need to know which medicine caused a symptom before calling.

1 Head / thyroid / energy

Fruquintinib

Fatigue and thyroid changes can affect daily life. Severe headache, confusion, seizure or visual change needs emergency help.

3 Heart / blood pressure

Fruquintinib

Blood pressure may rise without symptoms. Keep scheduled checks. Chest pain, collapse or severe breathlessness needs emergency care.

5 Liver


Fruquintinib

Liver tests can change before symptoms occur. Report yellow eyes or skin, dark urine or persistent abdominal pain.

7 Hands / feet / skin

Fruquintinib

Pain, redness, swelling or blisters can affect walking or using your hands. Report early and follow the skin-care plan.



2 Mouth / appetite

Fruquintinib

Mouth soreness, voice change and poor appetite can affect eating. Report difficulty drinking or weight loss.

4 Blood / bleeding

Fruquintinib

New bleeding, easy bruising, fever or sudden illness needs prompt advice. Heavy bleeding needs emergency help.

6 Bowel / abdomen

Fruquintinib

Diarrhoea can cause dehydration. Severe persistent pain, black stools or blood in stools needs urgent help.

8 Kidneys / urine

Fruquintinib

Protein in urine may have no early symptoms. Attend urine checks and report swelling or much less urine.

Selected effects, not every possible effect or a prediction. Blood-count, skin and line markers are schematic. Number and text carry meaning; colour is decorative.

Get urgent help for severe or persistent tummy pain, black or bloody stools, coughing or vomiting blood, new severe headache, confusion, seizure, visual change, chest pain, sudden breathlessness or very little urine. Report fever, sudden illness, painful hands/feet and new bleeding promptly.

When to get help

Call 112 / 999 for severe breathing difficulty, severe chest pain, collapse or a seizure.

Call oncology immediately: temperature 37.5°C or above, 36°C or below, shivering or suddenly feeling unwell, even with a normal temperature. Follow your hospital's temperature instructions. If you cannot reach the team, seek urgent medical assessment. Do not wait for the next appointment.

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.

Keep your treatment details to hand

Metastatic colorectal cancer after standard therapies and trifluridine/tipiracil or regorafenib · NCCP 00890 v2a · Document revised 5 November 2026

Print this page **one-sided on A4 at Actual size / 100%**. Cut only the dashed outer border around both panels. **Do not cut the centre line.** Fold with FRONT and BACK facing out, then glue or tape the blank backs together. Finished size: **85.6 × 54 mm**. Keep the stamp side writable.

FRONT / CONTACTS

My Treatment Card

SACTCheck

Name: _____

Fruquintinib (Fruzaqla)

28 days · capsules Days 1–21 · no capsules Days 22–28.

Phase: _____ Cycle plan: team to confirm

Daytime (EXAMPLE) 087 12345678

Out of hours (EXAMPLE) 087 87654321

Unwell or fever? Call oncology now.
Severe breathing difficulty, chest pain or collapse: **112 / 999.**

EXAMPLE NUMBERS — replace before patient use.

BACK / TREATMENT RECORD

My cycle record

Rolling record — follow your agreed treatment plan.

Cycle	Date / initials

My treatment information

Latest treatment date: _____

Plan changed? Ask the team to update this card.

CUT outer border

FOLD centre — do not cut

Keep the record accurate

The team dates and initials each recorded treatment unit when complete. A cycle is complete after its final planned dose; a weekly-treatment stamp records that weekly dose. Delays, omitted doses or a changed plan need a note or a replacement card. A stamp does not approve the next treatment. Keep any medicine-specific alert card too.

Put your treatment page on your phone

Scan your QR and check the treatment name. Save that page to your home screen.

iPhone / Safari

Tap Share → Add to Home Screen. Select Open as Web App if offered, then Add.

Android / Chrome

Tap the three-dot menu → Add to Home screen or Install and create shortcut. Choose the shortcut option and confirm.

Menu wording varies. The QR opens information, not a personal record. Paper stamps do not update the website. Check that the saved icon opens the correct treatment.

Your local contacts and resources

Oncology daytime: 087 12345678 · **Out of hours:** 087 87654321

EXAMPLE ONLY — team must replace both numbers before patient use.

Hospital / team: _____

Backup if the team cannot be reached: _____

Additional local services and resources will be added when confirmed. Your team must check and complete this page before patient use.

Feasibility build. Clinical, pharmacy and patient/PPI review pending. Your team confirms your personal plan.