

Modified FOLFOX-6

Three medicines · one 14-day cycle · a take-home pump

NCCP regimen 00209 · version 10a · colorectal / upper GI indications

WHAT IS IN THIS TREATMENT?

OXALIPLATIN

Platinum chemotherapy

FOLINIC ACID

Helps 5-FU work

5-FLUOROURACIL (5-FU)

Chemotherapy: injection + pump

YOUR CYCLE / REPEATS EVERY 14 DAYS

DAY 1

Hospital: oxaliplatin + folinic acid, then 5-FU. Pump starts.

DAY 2-3

Pump runs for 46 hours; disconnect is usually day 3.

DAY 3-6

Cold sensitivity can start early. Nausea, bowel or mouth changes may develop.

DAY 7-14

Blood counts may be lower; infection can occur on ANY day.

DAY 15

Next cycle if review and bloods allow.

These are possible time windows, not a forecast. A new symptom matters whenever it occurs; do not wait for a marked day.

BEFORE EACH CYCLE AND AT HOME

Before first 5-FU, DPD status is checked. Before each cycle: blood count, kidney/liver bloods and nerve symptoms. Report diarrhoea, mouth sores and painful red hands or feet.

My treatment purpose: _____ Planned cycles / review: _____

SUPPORT MEDICINES / FOLLOW YOUR OWN LABELLED PRESCRIPTION

CUH moderate-risk proforma: dexamethasone days 2-3; omeprazole days 1-3; nystatin and chlorhexidine mouth care; cyclizine and metoclopramide as prescribed for nausea; loperamide as prescribed for diarrhoea. Your actual list, doses and changes come from your oncology team.

Keep the pump dry and secure. If it leaks, disconnects or alarms, follow the pump instructions and contact the day ward.

CALL YOUR ONCOLOGY SOS LINE NOW


Temperature $\geq 37.5^{\circ}\text{C}$ or $\leq 36^{\circ}\text{C}$; shivers or feeling suddenly unwell; 4+ diarrhoea episodes in a day; vomiting despite medicine; mouth sores stopping fluids; bleeding; new chest pain or breathlessness.

Measure your temperature BEFORE paracetamol or ibuprofen if you feel unwell. Ask oncology for advice before using fever-reducing medicine; it can hide fever. Severe chest pain or breathing difficulty: call 112/999.

Your 24-hour oncology number: _____

Where FOLFOX can affect you

Oxaliplatin = cold + nerves 5-FU = gut + mouth + blood + rarely heart



COLD PRECAUTIONS + IMPORTANT PRACTICAL POINTS

For several days after oxaliplatin, avoid cold drinks, ice, the freezer and cold air if these trigger symptoms. Do not use ice chips during 5-FU bolus. New or worsening symptoms should be reported; do not diagnose a tight throat yourself.


Fertility, pregnancy and contraception advice is individual: ask your treating team before treatment.

READ MORE / TREATMENT RESOURCES

The companion extended portal contains detailed side-effect advice, source links and a printable treatment passport.

Scan to open the full FOLFOX patient portal on sactcheck.com. No patient details are in the QR.

Official regimen + HPRA product links in the portal



Full patient portal