NCCP 00448 · clinician draft

Panitumumab + FOLFIRI

Supportive prescription draft for review. This is a regimen template. It is not an issued patient prescription; individual review and completion are required.

Panitumumab + FOLFIRI

Active medicines: panitumumab + irinotecan + folinic acid + fluorouracil

Emetogenic risk: moderate

Medicine	Dose / route	When
Ondansetron	16 mg orally	Day 1 before chemotherapy Dose may be adjusted for individual/QT risk.
Dexamethasone	8 mg orally daily	Days 1–3 Reconcile all steroid prescriptions.
Metoclopramide	10 mg orally up to three times daily as prescribed	As required on subsequent days Confirm suitability, duration and contraindications.
Atropine	250 micrograms subcutaneously	Before irinotecan per NCCP prophylaxis Check contraindications including glaucoma; this does not treat delayed diarrhoea.

Additional phase-specific support

- No routine panitumumab premedication unless previous hypersensitivity/local policy indicates.
- Written delayed-diarrhoea plan and urgent fever/dehydration advice; do not confuse with acute cholinergic management.
- Skin care and electrolytes; monthly Mg/Ca/electrolytes for two months after last panitumumab.
- No routine panitumumab premedication unless previous hypersensitivity/local policy indicates. Written delayed-diarrhoea plan and urgent fever/dehydration advice; do not confuse with acute cholinergic management. Skin care and electrolytes; monthly Mg/Ca/electrolytes for two months after last panitumumab.

Before signing: verify allergies, current medicines, interactions, renal / hepatic function, QT risk, sedation, prior response, total steroid exposure and the rescue-medicine duration. Do not duplicate medicines from another phase.

Individual changes / rescue duration: _____
Prescriber / date: _____ Pharmacy review: _____

NCCP 00448 v5 · p6 · NCCP NCIS Medical Oncology antiemetics V8, Table 1

Sources checked 2026-11-04 · SACTCheck v0.84.21 · Regimen-level drafts; prescriber and oncology-pharmacy verification required