SACTCHECK · REGIMEN-SPECIFIC SACT CONSENT

Bevacizumab + FOLFOXIRI

Metastatic colorectal cancer · NCCP 00783 · version 2a · 14-day cycle

PROTOTYPE · GOVERNANCE REVIEW REQUIRED

Patient name	DOB	Hospital no.	Date

Treatment context
This combination treats colorectal cancer that has spread. Three chemotherapy medicines act on cancer cells while bevacizumab blocks a signal supporting new blood vessels. Treatment aims and any possibility of later local treatment are discussed individually.

Schedule
Day 1: bevacizumab, irinotecan, oxaliplatin and folinic acid, followed by a 48-hour fluorouracil pump. Repeat every 14 days while beneficial and tolerable.

1. TREATMENT DISCUSSION

- ☐ Purpose, expected benefit and alternatives explained
- ☐ Indication, medicines, sequence and duration confirmed
- ☐ Blood tests and any individually indicated heart checks discussed
- ☐ DPD result and severe fluoropyrimidine risks reviewed
- ☐ Support medicines and phase changes explained
- ☐ Fertility, contraception and breastfeeding discussed where relevant
- ☐ Common, serious and lasting drug-specific risks discussed below
- ☐ Urgent symptoms and local contact details confirmed

2. REGIMEN / AGENT-SPECIFIC MATERIAL RISKS

Bevacizumab
ANTI-VEGF TARGETED ANTIBODY

- ☐ High blood pressure / urine protein
- ☐ Bleeding or blood clots
- ☐ Delayed wound healing
- ☐ Rare bowel perforation / fistula
- ☐ Rare serious neurological effects
- ☐ Infusion reaction

Irinotecan
TOPOISOMERASE I CHEMOTHERAPY

- ☐ Early cramps, sweating and diarrhoea
- ☐ Delayed diarrhoea and dehydration
- ☐ Low blood counts / serious infection
- ☐ Nausea, mouth soreness and fatigue
- ☐ Hair thinning or loss
- ☐ Rare serious lung or allergic reactions

Oxaliplatin
PLATINUM CHEMOTHERAPY

- ☐ Cold-triggered tingling / throat discomfort
- ☐ Persistent sensory neuropathy, sometimes lasting after treatment
- ☐ Allergic / infusion reaction
- ☐ Low blood counts and infection
- ☐ Nausea, diarrhoea and fatigue
- ☐ Rare serious lung or blood complications

Folinic acid
CHEMOTHERAPY MODULATOR

- ☐ Enhances fluorouracil activity and toxicity
- ☐ Rare allergic reaction
- ☐ Diarrhoea / mouth soreness in combination

Fluorouracil
CHEMOTHERAPY

- ☐ Diarrhoea, dehydration and mouth ulcers
- ☐ Low blood counts / infection
- ☐ Hand-foot skin soreness
- ☐ Cardiotoxicity including chest pain / ischaemia
- ☐ Severe early toxicity including DPD-related toxicity
- ☐ Pump / central-line complications

Confirm the individual aim, likely benefit and alternatives, DPD result, the 48-hour pump, disconnection and urgent contacts. Explain early irinotecan symptoms separately from delayed diarrhoea. Explain cold sensitivity and potentially persistent neuropathy. Explain blood-pressure/urine checks, bleeding/clot risks and procedure/wound precautions.

Individual aim / benefit, alternatives, material risks and questions:

3. CONSENT CONFIRMATION

- ☐ Expected benefit, important risks and reasonable alternatives discussed
- ☐ Patient questions invited and answered; patient-specific material risks / priorities considered
- ☐ Emergency contact / toxicity advice and written information offered where appropriate
- ☐ Patient understands consent is voluntary and may be withdrawn

Patient / person giving consent _____ Date _____

Clinician obtaining consent _____ Date _____

Prototype only. This one-page form is a regimen-specific consent aid and requires formal governance, consultant and oncology-pharmacy review before clinical use. The current NCCP regimen, local/national consent policy, current medicine information, patient-specific material risks and clinical judgement remain authoritative.
Base regimen: NCCP 00783 v2a.

SACTCheck v0.84.33
Developed by Dr Paul O'Brien